


TITLE

FATTY ACID DESATURASE GENES FROM PLANTS

FIELD OF THE INVENTION

INS A' 

5 The invention relates to the preparation and use of nucleic acid fragments encoding fatty acid desaturase enzymes to modify plant lipid composition.

BACKGROUND OF THE INVENTION

Plant lipids have a variety of industrial and nutritional uses and are central to plant membrane function and climatic adaptation. These lipids represent a vast array of chemical structures, and these structures determine the physiological and industrial properties of the lipid. Many of these structures result either directly or indirectly from metabolic processes that alter the degree of unsaturation of the lipid. Different metabolic regimes in different plants produce these altered lipids, and either domestication of exotic plant species or modification of agronomically adapted species is usually required to economically produce large amounts of the desired lipid.

10
15
20

Plant lipids find their major use as edible oils in the form of triacylglycerols. The specific performance and health attributes of edible oils are determined largely by their fatty acid composition. Most vegetable oils derived from commercial plant varieties are composed primarily of palmitic (16:0), stearic (18:0), oleic (18:1), linoleic (18:2) and linolenic (18:3) acids. Palmitic and stearic acids are, respectively, 16- and 18-carbon-long, saturated fatty acids. Oleic, linoleic, and linolenic acids are 18-carbon-long, unsaturated fatty acids containing one, two, and three double bonds, respectively. Oleic acid is referred to as a mono-unsaturated fatty acid, while linoleic and linolenic acids are referred to as poly-unsaturated fatty acids. The relative amounts of saturated and

25
30
35

0049315667260

unsaturated fatty acids in commonly used, edible vegetable oils are summarized below (Table 1):

TABLE 1

Percentages of Saturated and Unsaturated Fatty Acids in the Oils of Selected Oil Crops

	<u>Saturated</u>	<u>Mono-unsaturated</u>	<u>Poly-unsaturated</u>
<u>Canola</u>	6%	58%	36%
<u>Soybean</u>	15%	24%	61%
<u>Corn</u>	13%	25%	62%
<u>Peanut</u>	18%	48%	34%
<u>Safflower</u>	9%	13%	78%
<u>Sunflower</u>	9%	41%	51%
<u>Cotton</u>	30%	19%	51%

Many recent research efforts have examined the role that saturated and unsaturated fatty acids play in reducing the risk of coronary heart disease. In the past, it was believed that mono-unsaturates, in contrast to saturates and poly-unsaturates, had no effect on serum cholesterol and coronary heart disease risk. Several recent human clinical studies suggest that diets high in mono-unsaturated fat and low in saturated fat may reduce the "bad" (low-density lipoprotein) cholesterol while maintaining the "good" (high-density lipoprotein) cholesterol (Mattson et al., Journal of Lipid Research (1985) 26:194-202).

A vegetable oil low in total saturates and high in mono-unsaturates would provide significant health benefits to consumers as well as economic benefits to oil processors. As an example, canola oil is considered a very healthy oil. However, in use, the high level of poly-unsaturated fatty acids in canola oil renders the oil unstable, easily oxidized, and susceptible to development of disagreeable odors and flavors (Gailliard, 1980, Vol. 4, pp. 85-116 In: Stumpf, P. K., Ed., The Biochemistry of Plants, Academic Press, New

York). The levels of poly-unsaturates may be reduced by hydrogenation, but the expense of this process and the concomitant production of nutritionally questionable trans isomers of the remaining unsaturated fatty acids reduces the overall desirability of the hydrogenated oil (Mensink et al., New England J. Medicine (1990) N323: 439-445). Similar problems exist with soybean and corn oils.

For specialized uses, high levels of poly-unsaturates can be desirable. Linoleate and linolenate are essential fatty acids in human diets, and an edible oil high in these fatty acids can be used for nutritional supplements, for example in baby foods. Linseed oil, derived from the Flax plant (*Linum usitatissimum*), contains over 50% linolenic acid and has widespread use in domestic and industrial coatings since the double bonds of the fatty acids react rapidly with oxygen to polymerize into a soft and flexible film. Although the oil content of flax is comparable to canola (around 40% dry weight of seed), high yields are only obtained in warm temperatures or subtropical climates. In the USA flax is highly susceptible to rust infection. It will be commercially useful if a crop such as soybean or canola could be genetically transformed by the appropriate desaturase gene(s) to synthesize oils with a high linolenic acid content.

Mutation-breeding programs have met with some success in altering the levels of poly-unsaturated fatty acid levels found in the edible oils of agronomic species. Examples of commercially grown varieties are high (85%) oleic sunflower and low (2%) linolenic flax (Knowles, (1980) pp. 35-38 In: Applewhite, T. H., Ed., World Conference on Biotechnology for the Fats and Oils Industry Proceedings, American Oil Chemists' Society). Similar commercial progress with the other plants shown

2025 RELEASE UNDER E.O. 14176

in Table 1 has been largely elusive due to the difficult nature of the procedure and the pleiotropic effects of the mutational regime on plant hardiness and yield potential.

5 The biosynthesis of the major plant lipids has been the focus of much research (Browse et al., Ann. Rev. Plant Physiol. Mol. Biol. (1991) 42:467-506). These studies show that, with the notable exception of the soluble stearyl-acyl carrier protein desaturase, the
10 controlling steps in the production of unsaturated fatty acids are largely catalyzed by membrane-associated fatty acid desaturases. Desaturation reactions occur in plastids and in the endoplasmic reticulum using a variety of substrates including galactolipids,
15 sulfolipids, and phospholipids. Genetic and physiological analyses of Arabidopsis thaliana nuclear mutants defective in various fatty acid desaturation reactions indicates that most of these reactions are catalyzed by enzymes encoded at single genetic loci in
20 the plant. The analyses show further that the different defects in fatty acid desaturation can have profound and different effects on the ultra-structural morphology, cold sensitivity, and photosynthetic capacity of the plants (Ohlrogge, et al., Biochim. Biophys. Acta (1991)
25 1082:1-26). However, biochemical characterization of the desaturase reactions has been meager. The instability of the enzymes and the intractability of their proper assay has largely limited researchers to investigations of enzyme activities in crude membrane
30 preparations. These investigations have, however, demonstrated the role of delta-12 desaturase and delta-15 desaturase activities in the production of linoleate and linolenate from 2-oleoyl-phosphatidylcholine and 2-linoleoyl-phosphatidylcholine,
35 respectively (Wang et al., Plant Physiol. Biochem.

00019935-122398
862221-5661260

5 Genes from plants for stearoyl-acyl carrier protein
desaturase, the only soluble fatty acid desaturase
known, have been described (Thompson, et al., Proc.
Nat'l. Acad. Sci. U.S.A. (1991) 88:2578-2582; Shanklin et
al., Proc. Nat'l. Acad. Sci. USA (1991) 88:2510-2514).
10 Stearoyl-coenzyme-A desaturase genes from yeast, rat,
and mice have also been described (Stukey, et al., J.
Biol. Chem. (1990) 265:20144-20149; Thiede, et al., J.
Biol. Chem. (1986) 261:13230-13235; Kaestner, et al., J.
Biol. Chem. (1989) 264:14755-1476). No evidence exists
15 in the public art that describes the isolation of fatty
acid desaturases other than stearoyl-ACP desaturases
from higher plants or their corresponding genes. A
fatty acid desaturase gene from the cyanobacterium,
Synechocystis PCC 6803, has also been described (Wada,
20 et al., Nature (1990) 347:200-203). This gene encodes a
fatty acid desaturase, designated des A, that catalyzes
the conversion of oleic acid at the 1 position of
galactolipids to linoleic acid. However, these genes
have not proven useful for isolating plant fatty acid
25 desaturases other than stearoyl-ACP desaturase via
sequence-dependent protocols, and the present art does
not indicate how to obtain plant fatty acid desaturases
other than stearoyl-ACP desaturases or how to obtain
fatty acid desaturase-related enzymes. Thus, the
30 present art does not teach how to obtain glycerolipid
desaturases from plants. Furthermore, there is no
evidence that a method to control the nature and levels
of unsaturated fatty acids in plants using nucleic acids
encoding fatty acid desaturases other than stearoyl-ACP
35 desaturase is known in the art.

The biosynthesis of the minor plant lipids has been less well studied. While hundreds of different fatty acids have been found, many from the plant kingdom, only a tiny fraction of all plants have been surveyed for their lipid content (Gunstone, et al., Eds., (1986) The Lipids Handbook, Chapman and Hall Ltd., Cambridge). Accordingly, little is known about the biosynthesis of these unusual fatty acids and fatty acid derivatives. Interesting chemical features found in such fatty acids include, for example, allenic and conjugated double bonds, acetylenic bonds, trans double bonds, multiple double bonds, and single double bonds in a wide number of positions and configurations along the fatty acid chain. Similarly, many of the structural modifications found in unusual lipids (e.g., hydroxylation, epoxidation, cyclization, etc.) are probably produced via further metabolism following chemical activation of the fatty acid by desaturation or they involve a chemical reaction that is mechanistically similar to desaturation. For example, evidence for the mechanism of hydroxylation of fatty acids being part of a general mechanism of enzyme-catalyzed desaturation in eukaryotes has been obtained by substituting a sulfur atom in the place of carbon at the delta-9 position of stearic acid. When incubated with yeast cell extracts the thiostearate was converted to a 9-sulfoxide (Buist et al. (1987) Tetrahedron Letters 28:857-860). This sulfoxidation was specific for sulfur at the delta-9 position and did not occur in a yeast delta-9-desaturase deficient mutant (Buist & Marecak (1991) Tetrahedron Letters 32:891-894). The 9-sulfoxide is the sulfur analogue of 9-hydroxyoctadecastearate, the proposed intermediate of stearate desaturation. Thus fatty-acid desaturase cDNAs may serve as useful probes for cDNAs encoding fatty-acid hydroxylases and other cDNAs which encode enzymes with

5 induced to synthesize them by introduction of a gene
encoding the appropriate desaturase.

SUMMARY OF THE INVENTION

Applicants have discovered a means to control the nature and levels of unsaturated fatty acids in plants. Nucleic acid fragments from glycerolipid desaturase cDNAs or genes are used to create chimeric genes. The chimeric genes may be used to transform various plants to modify the fatty acid composition of the plant or the oil produced by the plant. More specifically, one embodiment of the invention is an isolated nucleic acid fragment comprising a nucleotide sequence encoding a plant delta-15 fatty acid desaturase or a fatty acid desaturase-related enzyme with an amino acid identity of 50%, 65%, 90% or greater to the polypeptide encoded by SEQ ID NOS:1, 4, 6, 8, 10, 12, 14, or 16. The isolated fragment in these embodiments is isolated from a plant selected from the group consisting of soybean, oilseed Brassica species, Arabidopsis thaliana and corn.

Another embodiment of this invention involves the use of these nucleic acid fragments in sequence-dependent protocols. Examples include use of the fragments as hybridization probes to isolate other glycerolipid desaturase cDNAs or genes. A related embodiment involves using the disclosed sequences for amplification of DNA fragments encoding other glycerolipid desaturases.

Another aspect of this invention involves chimeric genes capable of causing altered levels of the linolenic acid in a transformed plant cell, the gene comprising
35 nucleic acid fragments encoding encoding a plant

delta-15 fatty acid desaturase or a fatty acid desaturase-related enzyme with an amino acid identity of 50%, 65%, 90% or greater to the polypeptide encoded by SEQ ID NOS:1, 4, 6, 8, 10, 12, 14, or 16 operably linked in suitable orientation to suitable regulatory sequences. Preferred are those chimeric genes which incorporate nucleic acid fragments encoding delta-15 fatty acid desaturase cDNAs or genes. Plants and oil from seeds of plants containing the chimeric genes described are also claimed.

Yet another embodiment of the invention involves a method of producing seed oil containing altered levels of linolenic (18:3) acid comprising: (a) transforming a plant cell with a chimeric gene described above; (b) growing fertile plants from the transformed plant cells of step (a); (c) screening progeny seeds from the fertile plants of step (b) for the desired levels of linolenic (18:3) acid, and (d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of the unsaturated fatty acids. Preferred plant cells and oils are derived from soybean, rapeseed, sunflower, cotton, cocoa, peanut, safflower, coconut, flax, oil palm, and corn. Preferred methods of transforming such plant cells would include the use of Ti and Ri plasmids of Agrobacterium, electroporation, and high-velocity ballistic bombardment.

The invention also is embodied in a method of breeding plant species to obtain altered levels of polyunsaturated fatty acids, specifically linolenic (18:3) acid in seed oil of oil-producing plants. This method involves (a) making a cross between two varieties of an oilseed plant differing in the linolenic acid trait; (b) making a Southern blot of restriction enzyme digested genomic DNA isolated from several progeny plants resulting from the cross of step (a); and (c)

hybridizing the Southern blot with the radiolabeled nucleic acid fragments encoding the claimed glycerolipid desaturases.

5 The invention is also embodied in a method of RFLP mapping that uses the isolated Arabidopsis thaliana delta-15 desaturase sequences described herein.

10 The invention is also embodied in plants capable of producing altered levels of glycerolipid desaturase by virtue of containing the chimeric genes described herein. Further, the invention is embodied by seed oil obtained from such plants.

15 The invention is also embodied in a method of RFLP mapping in a genomic RFLP marker comprising (a) making a cross between two varieties of plants; (b) making a Southern blot of restriction enzyme digested genomic DNA isolated from several progeny plants resulting from the cross of step (a); and (c) hybridizing the Southern blot with a radiolabelled nucleic acid fragments of the claimed fragments.

20 The invention is also embodied in a method to isolate nucleic acid fragments encoding fatty acid desaturases and fatty acid desaturase-related enzymes, comprising (a) comparing SEQ ID NOS:2, 5, 7, 9, 11, 13, 15 and 17 with other fatty acid desaturase polypeptide sequences; (b) identifying the conserved sequence(s) of 25 4 or more amino acids obtained in step a; (c) making region-specific nucleotide probe(s) or oligomer(s) based on the conserved sequences identified in step b; and d) using the nucleotide probe(s) or oligomers(s) of step c 30 to isolate sequences encoding fatty acid desaturases and fatty-acid desaturase-related enzymes by sequence-dependent protocols. The product of the method of isolation method described is also part of the invention.

BRIEF DESCRIPTION OF THE SEQUENCE DESCRIPTIONS

The invention can be more fully understood from the following detailed description and the Sequence Descriptions which form a part of this application. The Sequence Descriptions contain the one letter code for nucleotide sequence characters and the three letter code for amino acids in conformity with the IUPAC-IUB standard described in Nucleic Acids Research 13:3021-3030 (1985) and 37 C.F.R. 1.822 which are incorporated herein by reference.

SEQ ID NO:1 shows the complete 5' to 3' nucleotide sequence of 1350 base pairs of the Arabidopsis cDNA which encodes delta-15 desaturase in plasmid pCF3. Nucleotides 46 to 48 are the putative initiation codon of the open reading frame (nucleotides 46 to 1206). Nucleotides 1204 to 1206 are the termination codon. Nucleotides 1 to 45 and 1207 to 1350 are the 5' and 3' untranslated nucleotides, respectively. The 386 amino acid protein sequence in SEQ ID NO:1 is that deduced from the open reading frame.

SEQ ID NO:2 is the deduced peptide of the open-reading frame of SEQ ID NO:1.

SEQ ID NO:3 is a partial nucleotide sequence of the Arabidopsis genomic DNA insert in plasmid pF1 which shows the genomic sequence in the region of the Arabidopsis genome that encodes delta-15 desaturase. Nucleotides 68-255 are identical to nucleotides 1-188 of SEQ ID NO:1. Nucleotides 47 to 49 and 56 to 58 are termination codons in the same reading frame as the open reading frame in SEQ ID NO:1.

SEQ ID NO:4 shows the 5' to 3' nucleotide sequence of the insert in plasmid pACF2-2 of 1525 base pairs of the Arabidopsis thaliana cDNA that encodes a plastid delta-15 fatty acid desaturase. Nucleotides 10-12 and nucleotides 1348 to 1350 are, respectively, the putative

5 SEQ ID NO:5 is the deduced peptide of the open
reading frame of SEQ ID NO:4.

SEQ ID NO:7 is the deduced peptide of the open reading frame of SEQ ID NO:6.

SEQ ID NO:9 is the deduced peptide of the open reading frame of SEQ ID NO:8.

SEQ ID NO:10 is the complete nucleotide sequence of the soybean (glycine max) microsomal delta-15 desaturase cDNA, found in plasmid pXF1, which the 2184 nucleotides of this sequence contain both the coding sequence and the 5' and 3' non-translated regions of the cDNA. Nucleotides 855 to 857 are the putative initiation codon of the open reading frame (nucleotides 855 to 2000). Nucleotides 1995 to 1997 are the termination codon.

5 SEQ ID NO:11 is the deduced peptide of the open
reading frame in SEQ ID NO:10.

20 SEQ ID NO:13 is the deduced peptide of the open
reading frame in SEQ ID NO:12.

SEQ ID NO:15 is the deduced amino acid sequence of
SEQ ID NO:14.

SEQ ID NO:16 shows the partial composite 5' to 3'
35 nucleotide sequence of 472 bp derived from the inserts

5

10

15

20

25

30

35

SEQ ID NO:26 Seventy two-fold degenerate antisense 18-mers made to amino acid residues 304-309 in SEQ ID NO:2.

5 SEQ ID NO:27 Seventy two-fold degenerate antisense 18-mers made to amino acid residues 304-309 in SEQ ID NO:2.

SEQ ID NO:28 Sixteen-fold degenerate sense 36-mers made to amino acid residues 97-108 in SEQ ID NO:2.

10 SEQ ID NO:29 Sixteen-fold degenerate sense 36-mers made to amino acid residues 97-108 in SEQ ID NO:2.

SEQ ID NO:30 Sixty four-fold degenerate antisense 38-mers made to amino acid residues 299-311 in SEQ ID NO:2.

15 SEQ ID NO:31 Sixty four-fold degenerate antisense 38-mers made to amino acid residues 299-311 in SEQ ID NO:2.

SEQ ID NO:32 A 135-mer made as an antisense strand to amino acid residues 97-141 in SEQ ID NO:2.

DETAILED DESCRIPTION OF THE INVENTION

20 Applicants have isolated nucleic acid fragments that encode plant fatty acid desaturases and that are useful in modifying fatty acid composition in oil-producing species by transformation.

25 Thus, transfer of the nucleic acid fragments of the invention or a part thereof that encodes a functional enzyme, along with suitable regulatory sequences that direct the transcription of their mRNA, into a living cell will result in the production or over-production of plant fatty acid desaturases and will result in
30 increased levels of unsaturated fatty acids in cellular lipids, including triacylglycerols.

Transfer of the nucleic acid fragments of the invention or a part thereof, along with suitable regulatory sequences that direct the transcription of
35 their antisense RNA, into plants will result in the

inhibition of expression of the endogenous fatty acid desaturase that is substantially homologous with the transferred nucleic acid fragment and will result in decreased levels of unsaturated fatty acids in cellular lipids, including triacylglycerols.

Transfer of the nucleic acid fragments of the invention or a part thereof, along with suitable regulatory sequences that direct the transcription of their mRNA, into plants may result in inhibition by cosuppression of the expression of the endogenous fatty acid desaturase gene that is substantially homologous with the transferred nucleic acid fragment and may result in decreased levels of unsaturated fatty acids in cellular lipids, including triacylglycerols.

The nucleic acid fragments of the invention can also be used as restriction fragment length polymorphism (RFLP) markers in Arabidopsis genetic mapping and plant breeding programs.

The nucleic acid fragments of the invention or oligomers derived therefrom can also be used to isolate other related glycerolipid desaturase genes using DNA, RNA, or a library of cloned nucleotide sequences from the same or different species by well known sequence-dependent protocols, including, for example, methods of nucleic acid hybridization and amplification by the polymerase chain reaction.

Definitions

In the context of this disclosure, a number of terms shall be used. The term "fatty acid desaturase" used herein refers to an enzyme which catalyzes the breakage of a carbon-hydrogen bond and the introduction of a carbon-carbon double bond into a fatty acid molecule. The fatty acid may be free or esterified to another molecule including, but not limited to, acyl-carrier protein, coenzyme A, sterols and the glycerol

moiety of glycerolipids. The term "glycerolipid desaturases" used herein refers to a subset of the fatty acid desaturases that act on fatty acyl moieties esterified to a glycerol backbone. "Delta-12

5 desaturase" refers to a fatty acid desaturase that catalyzes the formation of a double bond between carbon positions 6 and 7 (numbered from the methyl end), (i.e., those that correspond to carbon positions 12 and 13 (numbered from the carbonyl carbon) of an 18 carbon-long

10 fatty acyl chain or carbon positions 10 and 11 (numbered from the carbonyl carbon) of a 16 carbon-long fatty acyl chain). "Delta-15 desaturase" refers to a fatty acid desaturase that catalyzes the formation of a double bond between carbon positions 3 and 4 (numbered from the

15 methyl end), (i.e., those that correspond to carbon positions 15 and 16 (numbered from the carbonyl carbon) of an 18 carbon-long fatty acyl chain and carbon positions 13 and 14 (numbered from the carbonyl carbon) of a 16 carbon-long fatty acyl chain). Examples of

20 fatty acid desaturases include, but are not limited to, the microsomal delta-12 and delta-15 desaturases that act on phosphatidylcholine lipid substrates; the chloroplastic delta-12 and delta-15 desaturases that act on phosphatidyl glycerol and galactolipids; and other

25 desaturases that act on such fatty acid substrates such as phospholipids, galactolipids, and sulfolipids. "Microsomal desaturase" refers to the cytoplasmic location of the enzyme, while "chloroplast desaturase" and "plastid desaturase" refer to the plastid location

30 of the enzyme. These fatty acid desaturases may be found in a variety of organisms including, but not limited to, higher plants, diatoms, and various eukaryotic and prokaryotic microorganisms such as fungi and photosynthetic bacteria and algae. The term

35 "homologous fatty acid desaturases" refers to fatty acid

desaturases that catalyze the same desaturation on the same lipid substrate. Thus, microsomal delta-15 desaturases, even from different plant species, are homologous fatty acid desaturases. The term

5 "heterologous fatty acid desaturases" refers to fatty acid desaturases that catalyze desaturations at different positions and/or on different lipid substrates. Thus, for example, microsomal delta-12 and delta-15 desaturases, which act on phosphatidylcholine

10 lipids, are heterologous fatty acid desaturases, even when from the same plant. Similarly, microsomal delta-15 desaturase, which acts on phosphatidylcholine lipids, and chloroplast delta-15 desaturase, which acts on galactolipids, are heterologous fatty acid

15 desaturases, even when from the same plant. It should be noted that these fatty acid desaturases have never been isolated and characterized as proteins. Accordingly the terms such as "delta-12 desaturase" and "delta-15 desaturase" are used as a convenience to

20 describe the proteins encoded by nucleic acid fragments that have been isolated based on the phenotypic effects caused by their disruption. The term "fatty acid desaturase-related enzyme" refers to enzymes whose catalytic product may not be a carbon-carbon double bond

25 but whose mechanism of action is similar to that of a fatty acid desaturase (that is, catalysis of the displacement of a carbon-hydrogen bond of a fatty acid chain to form a fatty-hydroxyacyl intermediate or end-product). This term is different from "related fatty

30 acid desaturases", which refers to structural similarities between fatty acid desaturases.

The term "nucleic acid" refers to a large molecule which can be single-stranded or double-stranded, composed of monomers (nucleotides) containing a sugar, a

35 phosphate and either a purine or pyrimidine. A "nucleic

acid fragment" is a fraction of a given nucleic acid molecule. In higher plants, deoxyribonucleic acid (DNA) is the genetic material while ribonucleic acid (RNA) is involved in the transfer of the information in DNA into proteins. A "genome" is the entire body of genetic material contained in each cell of an organism. The term "nucleotide sequence" refers to the sequence of DNA or RNA polymers, which can be single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases capable of incorporation into DNA or RNA polymers. The term "oligomer" refers to short nucleotide sequences, usually up to 150 bases long. "Region-specific nucleotide probes" refers to isolated nucleic acid fragments derived from a cDNA or gene using a knowledge of the amino acid regions conserved between different fatty-acid desaturases which may be used to isolate cDNAs or genes for other fatty-acid desaturases or fatty acid desaturase-related enzymes using sequence dependent protocols. As used herein, the term "homologous to" refers to the relatedness between the nucleotide sequence of two nucleic acid molecules or between the amino acid sequences of two protein molecules. Estimates of such homology are provided by either DNA-DNA or DNA-RNA hybridization under conditions of stringency as is well understood by those skilled in the art (Hames and Higgins, Eds. (1985) Nucleic Acid Hybridisation, IRL Press, Oxford, U.K.); or by the comparison of sequence similarity between two nucleic acids or proteins, such as by the method of Needleman et al. (J. Mol. Biol. (1970) 48:443-453). As used herein, "substantially homologous" refers to nucleotide sequences that have more than 90% overall identity at the nucleotide level with the coding region of the claimed sequence, such as genes and pseudo-genes corresponding to the coding

091936-1239

regions. The nucleic acid fragments described herein include molecules which comprise possible variations, both man-made and natural, such as but not limited to (a) those that involve base changes that do not cause a change in an encoded amino acid, or (b) which involve base changes that alter an amino acid but do not affect the functional properties of the protein encoded by the DNA sequence, (c) those derived from deletions, rearrangements, amplifications, random or controlled mutagenesis of the nucleic acid fragment, and (d) even occasional nucleotide sequencing errors.

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding) and following (3' non-coding) the coding region. "Fatty acid desaturase gene" refers to a nucleic acid fragment that expresses a protein with fatty acid desaturase activity. "Native" gene refers to an isolated gene with its own regulatory sequences as found in nature. "Chimeric gene" refers to a gene that comprises heterogeneous regulatory and coding sequences not found in nature. "Endogenous" gene refers to the native gene normally found in its natural location in the genome and is not isolated. A "foreign" gene refers to a gene not normally found in the host organism but that is instead introduced by gene transfer. "Pseudo-gene" refers to a genomic nucleotide sequence that does not encode a functional enzyme.

"Coding sequence" refers to a DNA sequence that codes for a specific protein and excludes the non-coding sequences. It may constitute an "uninterrupted coding sequence", i.e., lacking an intron or it may include one or more introns bounded by appropriate splice junctions. An "intron" is a nucleotide sequence that is transcribed in the primary transcript but that is removed through cleavage and re-ligation of the RNA within the cell to

create the mature mRNA that can be translated into a protein.

"Initiation codon" and "termination codon" refer to a unit of three adjacent nucleotides in a coding sequence that specifies initiation and chain termination respectively, of protein synthesis (mRNA translation). "Open reading frame" refers to the coding sequence uninterrupted by introns between initiation and termination codons that encodes an amino acid sequence.

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA.

"Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to RNA transcript that includes the mRNA.

"Antisense RNA" refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene by interfering with the processing, transport and/or translation of its primary transcript or mRNA. The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. In addition, as used herein, antisense RNA may contain regions of ribozyme sequences that increase the efficacy of antisense RNA to block gene expression. "Ribozyme" refers to a catalytic RNA and includes sequence-specific endoribonucleases.

SECRET 5657260

As used herein, "suitable regulatory sequences" refer to nucleotide sequences in native or chimeric genes that are located upstream (5'), within, and/or downstream (3') to the nucleic acid fragments of the invention, which control the expression of the nucleic acid fragments of the invention. The term "expression", as used herein, refers to the transcription and stable accumulation of the sense (mRNA) or the antisense RNA derived from the nucleic acid fragment(s) of the invention that, in conjunction with the protein apparatus of the cell, results in altered levels of the fatty acid desaturase(s). Expression or overexpression of the gene involves transcription of the gene and translation of the mRNA into precursor or mature fatty acid desaturase proteins. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of preventing the expression of the target protein. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. "Cosuppression" refers to the expression of a foreign gene which has substantial homology to an endogenous gene resulting in the suppression of expression of both the foreign and the endogenous gene. "Altered levels" refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

"Promoter" refers to a DNA sequence in a gene, usually upstream (5') to its coding sequence, which controls the expression of the coding sequence by providing the recognition for RNA polymerase and other factors required for proper transcription. In artificial DNA constructs promoters can also be used to transcribe antisense RNA. Promoters may also contain DNA sequences that are involved in the binding of

protein factors which control the effectiveness of transcription initiation in response to physiological or developmental conditions. It may also contain enhancer elements. An "enhancer" is a DNA sequence which can stimulate promoter activity. It may be an innate element of the promoter or a heterologous element inserted to enhance the level and/or tissue-specificity of a promoter. "Constitutive promoters" refers to those that direct gene expression in all tissues and at all times. "Tissue-specific" or "development-specific" promoters as referred to herein are those that direct gene expression almost exclusively in specific tissues, such as leaves or seeds, or at specific development stages in a tissue, such as in early or late embryogenesis, respectively.

The "3' non-coding sequences" refers to the DNA sequence portion of a gene that contains a polyadenylation signal and any other regulatory signal capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor.

The term "Transit Peptide" refers to the N-terminal extension of a protein that serves as a signal for uptake and transport of that protein into an organelle such as a plastid or mitochondrion.

"Transformation" herein refers to the transfer of a foreign gene into the genome of a host organism and its genetically stable inheritance. "Restriction fragment length polymorphism" refers to different sized restriction fragment lengths due to altered nucleotide sequences in or around variant forms of genes. "Fertile" refers to plants that are able to propagate sexually.

SECRET 5261340

5 (including Brassica napus, B. campestris), sunflower (Helianthus annuus), cotton (Gossypium hirsutum), corn (Zea mays), cocoa (Theobroma cacao), safflower (Carthamus tinctorius), oil palm (Elaeis guineensis), coconut palm (Cocos nucifera), flax (Linum

"Sequence-dependent protocols" refer to techniques that rely on a nucleotide sequence for their utility. Examples of sequence-dependent protocols include, but are not limited to, the methods of nucleic acid and oligomer hybridization and methods of DNA and RNA amplification such as are exemplified in various uses of the polymerase chain reaction. "PCR product" refers to the DNA product obtained through polymerase chain reaction.

Various solutions used in the experimental manipulations are referred to by their common names such as "SSC", "SSPE", "Denhardt's solution", etc. The composition of these solutions may be found by reference to Appendix B of Sambrook, et al. (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press).

T-DNA Mutagenesis and Identification of an
Arabidopsis Mutant Defective in Delta-15 Desaturation

In T-DNA mutagenesis (Feldmann, et al., Science (1989) 243:1351-1354), the integration of T-DNA in the genome can interrupt normal expression of the gene at or near the site of the integration. If the resultant mutant phenotype can be detected and shown genetically to be tightly linked to the T-DNA insertion, then the "tagged" locus and its wild type counterpart can be readily isolated by molecular cloning by one skilled in the art.

Arabidopsis thaliana seeds were transformed by Agrobacterium tumefaciens C58Clrif strain harboring the avirulent Ti-plasmid pGV3850::pAK1003 that has the T-DNA region between the left and right T-DNA borders replaced by the origin of replication region and ampicillin resistance gene of plasmid pBR322, a bacterial kanamycin resistance gene, and a plant kanamycin resistance gene (Feldmann, et al., Mol. Gen. Genetics (1987) 208:1-9). Plants from the treated seeds were self-fertilized and the resultant progeny seeds, germinated in the presence of kanamycin, were self-fertilized to give rise to a population, designated T3, that was segregating for T-DNA insertions. T3 seeds from approximately 6000 T2 plants were analyzed for fatty acid composition. One line, designated 3707, showed a reduced level of linolenic acid (18:3). One more round of self-fertilization of mutant line 3707 produced T4 progeny seeds. The ratio of 18:2/18:3 in seeds of the homogyzous mutant in T4 population was ca. 14; this ratio is ca 1.8 and ca. 23, respectively, in wild-type Arabidopsis and Arabidopsis fad 3 mutant [Lemieux et al. (1990) Theor. App. Gen. 80:234-240] obtained via chemical mutagenesis. These seeds were planted and individual plants were analyzed for the presence of

nopaline in leaf extracts. T5 seeds from these plants were further analyzed for fatty acid composition and the ability to germinate in the presence of kanamycin. The mutant fatty acid phenotype was found to segregate in a 1:2:1 ratio, as was germinability on kanamycin. Nopaline was found in all plants with an altered fatty acid phenotype, but not in wild type segregants. These results provided evidence that the locus controlling delta-15 desaturation was interrupted by T-DNA in mutant line 3707.

Isolation of Arabidopsis Genomic DNA

Containing the Gene Controlling Delta-15 Desaturation

In order to isolate the gene controlling delta-15 desaturation from wild-type Arabidopsis, a T-DNA-plant DNA "junction" fragment containing a T-DNA border integrated into the host plant DNA was isolated from Arabidopsis mutant 3707. For this, genomic DNA from the mutant plant was isolated and completely digested by either Bam HI or Sal I restriction enzymes. In each case, one of the resultant fragments was expected to contain the origin of replication and ampicillin-resistance gene of pBR322 as well as the left T-DNA-plant DNA junction fragment. Such fragments were rescued as plasmids by ligating the digested genomic DNA fragments at a dilute concentration to facilitate self-ligation and then using the ligated fragments to transform E. coli cells. Ampicillin-resistant E. coli transformants were isolated and screened by colony hybridization to fragments containing either the left or the right T-DNA border. Of the 192 colonies obtained from the plasmid rescue of Sal I digested genomic DNA, 31 hybridized with the left T-DNA border fragment, 4 hybridized to the right T-DNA border fragment, and none hybridized to both. Of the 85 colonies obtained from the plasmid rescue of Bam HI digested genomic DNA, 63

091936 1239
86227 5E6T260

hybridized to the left border and none to the right border. Restriction analysis of seven rescued plasmids that were obtained from the Bam HI digestion and that hybridized to the left T-DNA border showed that they
5 were indistinguishable and contained 1.4 kb of putative, flanking plant DNA. Restriction analysis of another rescued plasmid, pS1, that was obtained from the Sal I digestion and hybridized only to the left T-DNA border, showed that it contained 2.9 kb of putative, flanking
10 plant DNA. This flanking DNA had a Bam HI site and a Hind III site 1.4 kb and 2.2 kb, respectively, away from the left T-DNA border, suggesting that the 1.4 kb putative plant DNA in Bam HI rescued plasmids was contained within the 2.9 kb putative plant DNA in the
15 Sal I rescued plasmids. Southern blot analysis of wild type and mutant 3707 Arabidopsis genomic DNA using the radiolabeled 1.4 kb DNA fragment as the hybridization probe confirmed that this fragment contained plant DNA and that the T-DNA integration site was in a 2.8 kb
20 Bam HI, a 5.2 kb Hind III, a 3.5 kb Sal I, a 5.5 kb Eco RI, and an approximately 9 kb Cla I fragment of wild type Arabidopsis DNA. Nucleotide sequencing of plasmid pS1 with a primer made to a left T-DNA border sequence revealed that pS1 was colinear with the sequence of the
25 left T-DNA border (Yadav et al., Proc. Natl. Acad. Sci. USA (1982) 79:6322-6326) up to nucleotide position 65, which is in the T-DNA border repeats. Approximately 800 bp of additional sequence in pS1 beyond the T-DNA-plant DNA junction, that is, in the plant DNA adjoining the
30 left T-DNA border, showed no significant homology to the T-DNA of pGV3850::pAK1003 and no significant open reading frame.

The nucleic acid fragment from wild-type Arabidopsis corresponding to the plant DNA flanking
35 T-DNA in the line 3707 was isolated by screening a

lambda phage Arabidopsis thaliana genomic library with the 1.4 kb plant DNA isolated from the rescued plasmids as a hybridization probe. Seven positively-hybridizing genomic clones were isolated that fell in one of five classes based on partial restriction mapping. While their average insert size was approximately 15 kb, taken together they spanned a total of approximately 40 kb of genomic DNA. A combination of restriction and Southern analyses revealed that the five clones overlapped the site of integration of the left border of the T-DNA and that there was no detectable rearrangement of plant DNA in the rescued plasmids as compared to that in the wild type genomic plant DNA. One of these lambda phage clones, designated 1111, was representative of the recovered clones and contained an approximately 20 kb genomic DNA insert which was more or less symmetrically arranged around the site of insertion of the left border of the T-DNA. This clone was deposited on November 27, 1991 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 75167.

Isolation of Arabidopsis Delta-15

Desaturase cDNA

A 5.2 kb Hind III fragment containing wild-type genomic DNA, which hybridized to the 1.4 kb flanking plant DNA recovered from line 3707 and which was interrupted near its middle by the T-DNA insertion in line 3707, was isolated from lambda phage clone 41A1 and cloned into the Hind III site of the pBluescript SK vector (Stratagene) by standard cloning procedures described in Sambrook et al., Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press). The resultant plasmid was designated pF1. The isolated 5.2 kb Hind III fragment was also used as a radiolabeled hybridization probe to screen a

cDNA library made to poly A⁺ mRNA from 3-day-old etiolated Arabidopsis thaliana (ecotype Columbia) seedling hypocotyls in a lambda ZAP II vector (Stratagene). Of the several positively-hybridizing
5 plaques, four strongly-hybridizing ones were subjected to plaque purification. Sequences of the pBluescript (Stratagene) vector, including the cDNA inserts, from each of the purified phage stocks were excised in the presence of a helper phage. The resultant phagemids
10 were used to infect E. coli cells which yielded double-stranded plasmids, pCF1, pCF2, pCF3, and pCF4. All four were shown to contain at least one approximately 1.3 to 1.4 kb Not I insert fragment (Not I/Eco RI adaptors were used in the preparation of the cDNA library) which
15 hybridized to the same region of wild-type plant genomic DNA present in the isolated phage clones. This region, which was near the site of integration of the left T-DNA border in line 3707, was on the side of the T-DNA insertion opposite to that of the plant DNA flanking the
20 left T-DNA border isolated previously via plasmid rescue. Partial sequence determination of the different cDNAs revealed common identity. Since multiple versions of only one type of cDNA were obtained from a cDNA library made from etiolated tissue which is expected to
25 express delta-15 desaturation, and since these cDNAs hybridized to the genomic DNA that corresponds to the site of T-DNA integration in line 3707 which had a high linoleic acid/low linolenic acid phenotype, Applicants were lead to conclude that the T-DNA in line 3707
30 interrupted the normal expression of the gene encoding delta-15 desaturase. The complete nucleotide sequence of one cDNA, designated pCF3, was determined and is shown as SEQ ID NO:1. It reveals an open reading frame that encodes a 386 amino acid polypeptide. One of the
35 sequencing primers made to the pCF3 insert was also used

to obtain 255 bp of sequence from pF1 that is shown as SEQ ID NO:3. Nucleotides 68 to 255 of the genomic DNA in pF1 (SEQ ID NO:3) are identical to nucleotides 1 to 188 of the cDNA (SEQ ID NO:1), which shows that they are

5 colinear and that the cDNA is encoded for by the gene in the isolated genomic DNA. Nucleotides 113 to 115 in SEQ ID NO:3 are the initiation codon of the largest open reading frame corresponding to nucleotides 46-48 in SEQ ID NO:1. This is evident from the presence of in-frame

10 termination codons at nucleotides 47 to 49 and nucleotides 56 to 58 and the absence of observable intron splice junctions in SEQ ID NO:3. The identification of the 386 amino acid polypeptide as a

15 desaturase was confirmed by comparing its amino acid sequence with all the protein sequences found in Release 19.0 of the SWISSPROTEIN database using the FASTA algorithm of Pearson and Lipman (Proc. Natl. Acad. Sci. USA (1988) 85:2444-2448) and the BLAST program (Altschul et al., J. Mol. Biol. (1990) 215:403-410). The most

20 homologous protein found in both searches was the desA fatty acid desaturase from the cyanobacterium Synechocystis PCC6803 (Wada, et al., Nature (1990) 347:200-203; Genbank ID:CSDESA; GenBank Accession No:X53508). The 386 amino acid peptide in SEQ ID NO:1

25 was also compared to the 351 amino acid sequence of desA by the method of Needleman et al. (J. Mol. Biol. (1970) 48:443-453). Over their entire length, these proteins were 26% identical, the comparison imposing four major gaps in the desA protein sequence. While this overall

30 homology is poor, homology in shorter stretches was better. For instance, in a stretch of 78 amino acids the Arabidopsis delta-15 desaturase (amino acids 78 to 155 in SEQ ID NO:1) and the desA protein (amino acids 67 to 144) showed 40% identity and 66% similarity.

0930936
3663260

Homology in yet shorter stretches was even greater as shown in Table 2.

TABLE 2

Peptide Length	AA positions in SEQ ID NO:1	AA positions in desa	Percent Identity
12	97-108	86-97	83
7	115-121	104-110	71
9	133-141	22-130	56
11	299-309	282-292	64

These high percent identities in short stretches of amino acids between the cyanobacterial desaturase polypeptide and SEQ ID NO:2 suggests significant relatedness between the two.

To analyse the developmental expression of the gene encoding mRNA corresponding to SEQ ID NO:1, the cDNA insert in plasmid pCF3 was used as a radiolabeled hybridization probe on mRNA samples from leaf, root, germinating seedling, and developing siliques from both wild type and mutant 3707 Arabidopsis plants, essentially as described in Maniatis et al., Molecular Cloning, A Laboratory Manual (1982) Cold Spring Harbor Laboratory Press. The results indicated that while the mRNA corresponding to SEQ ID NO:1 is detected in all tissues from the mutant plant, its levels are lower than in wild-type tissues. This is consistent with the observation that the fatty acid mutation in line 3707 is leaky relative to the known Arabidopsis fad 3 mutant obtained via chemical mutagenesis. These results confirmed that the T-DNA in line 3707 had interrupted the normal expression of a fatty acid desaturase gene. Based on the fatty acid phenotype of homozygous mutant line 3707, Applicants concluded that the cDNA insert in pCF3 encoded the delta-15 desaturase. Further, Applicants concluded that it was the microsomal delta-15 desaturase, and not the chloroplastic delta-15

desaturase, since: a) the mutant phenotype was expressed strongly in the seed but expressed poorly, if at all, in the leaf of line 3707, and b) the delta-15 desaturase polypeptide, by comparison to the desA polypeptide, did not have an N-terminal extension of a transit peptide expected for a nuclear-encoded chloroplast desaturase.

The identity of SEQ ID NO:2 as the Arabidopsis microsomal delta-15 desaturase was confirmed by its biological overexpression in plant tissues. For this, the 1.4 kB Not I fragment of plasmid pCF3 containing the delta-15 desaturase cDNA was placed in the sense orientation behind either the CaMV 35S promoter, to provide constitutive expression, or behind the promoter for the gene encoding soybean a' subunit of the β -conglycinin (7S) seed storage protein, to provide embryo-specific expression. The chimeric genes 35S promoter/sense SEQ ID NO:1/3' nopaline synthase and β -conglycinin/sense SEQ ID NO:1/3' phaseolin were then transformed into plant cells by Agrobacterium tumefaciens's binary Ti plasmid vector system [Hoekema et al. (1983) Nature 303:179-180; Bevan (1984) Nucl. Acids Res. 12:8711-8720].

To confirm the identity of SEQ ID NO:1 and to test the biological effect of its overexpression in a heterologous plant species, the chimeric genes 35S promoter/sense SEQ ID NO:1/3' nopaline synthase was transformed into a binary vector, which was then transferred into Agrobacterium tumefaciens strain R1000, carrying the Ri plasmid pRiA4b from Agrobacterium rhizogenes [Moore et al. (1979) Plasmid 2:617-626]. Carrot (Daucus carota L.) cells were transformed by co-cultivation of carrot root disks with strain R1000 carrying the chimeric gene by the method of Petit et al. (1986) [Mol. Gen. Genet. 202:388-393]. Fatty acid

063227" 52667250

5 To complement the delta-15 desaturation mutation in
the T-DNA mutant line 3707 and to test the biological
effect of overexpression of SEQ ID NO:1 in seed, the
embryo-specific promoter/SEQ ID NO:1/3' phaseolin
chimeric gene was transformed into a binary vector,
0 which was then transformed into the avirulent
Agrobacterium strain LBA4404/pAL4404 [Hoekema et al.
(1983) Nature 303:179-180]. Roots of line 3707 were
transformed by the engineered Agrobacterium, transformed
plants were selected and grown to give rise to seeds.
5 Fatty acid analysis of the seeds from two plants showed
that the one out of six seeds in each plant showed the
mutant fatty acid phenotype, while the remaining seeds
show more than 10-fold increase in 18:3 to ca. 55%.
While the sample size is small, this segregation
0 suggests Mendelian inheritance of the fatty acid
phenotype. While most of the increase occurs at the
expense of 18:2, some of it also occurs at the expense
of 18:1. Thus, overexpression of this gene in oils
crops, especially canola, which is a close relative of
5 Arabidopsis, is also expected to result in the high
levels of 18:3 that are found in specialty oil of
linseed.

Comparisons of the sequence of the 386 amino acid polypeptide by the method of Needleman et al. (J. Mol. Biol. (1970) 48:443-453) with those for the microsomal stearyl-CoA (Δ^9) desaturases from rat, mouse and yeast revealed 21%, 19%, and 17% identities, respectively. While the membrane-associated Arabidopsis Δ^{15} desaturase protein showed significant but limited homology to the desA protein, it showed no

significant homology to the soluble stearoyl-ACP (delta-9) desaturases from higher plants, including one from Arabidopsis.

Comparison of partial nucleotide sequences of plasmids pF1 and pS1 showed that the left T-DNA border:plant DNA junction is ca. 700 bp from the initiation codon in SEQ ID NO:1. To determine the position of the other T-DNA:plant DNA junction with respect to the pF1 sequence, the T-DNA:plant DNA junction fragment was isolated. Genomic DNA from mutant line 3707, isolated as described previously, was partially digested by restriction enzyme Mbo I to give an average fragment size of ca. 15 kB. The fragment ends were partially-filled with dGTP and gATP by Klenow and cloned into Xho I half-sites of LambdaGEM[®]-11 (Promega Corporation) following the manufacturer's protocol. The phage library was titered and used essentially as described in Ausubel et al. [Current Protocols in Molecular Biology (1989) John Wiley & Sons]. The genomic phage library was screened with radiolabeled PCR product, ca. 0.6 kB, derived from 5' end of the gene in pF1. This product spans from 3 bp to the right of where the left-T-DNA border inserted to 15 bp to the left of nucleotide position 1 in SEQ ID NO:1. Southern blot analysis of DNA from one of the purified, positively-hybridizing phages following Eco RI restriction digestion and electrophoresis showed that a 4 kB Eco RI fragment hybridized to the 0.6 kB PCR product. The Eco RI fragment was subcloned and subject to sequence analyses. Comparison of the sequences derived from this fragment, pF1 and pS1 showed that the insertion of T-DNA resulted in a 56 bp deletion at the site of insertion and that the T-DNA interrupted the Arabidopsis gene 711 bp 5' to the initiation codon in SEQ ID NO:1. Thus, the T-DNA inserts 5' to the open reading

09019935-122350

frame, consistent with the leaky expresssion of the gene encoding SEQ ID NO:1 and the leaky fatty acid phenotype in mutant 3707. While the left T-DNA:plant DNA junction is precise, that is without any sequence rearrangement in either the left T-DNA border or the flanking plant DNA, the other T-DNA:plant DNA junction is complex and not fully characterized.

Plasmid pCF3 was deposited on December 3, 1991 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 68875.

Using Arabidopsis Delta-15 Desaturase cDNA as a
Hybridization Probe to Isolate cDNAs Encoding
Related Desaturases from Arabidopsis

The 1.4 kb Not I insert fragment isolated from plasmid pCF3 was purified, radiolabeled, and used to screen approximately 80,000 clones from the cDNA library made to poly A⁺ mRNA from 3-day-old etiolated Arabidopsis thaliana as described above, except that lower stringency hybridizations (1 M NaCl, 50 mM Tris-HCl, pH 7.5, 1% SDS, 5% dextran sulfate, 0.1 mg/mL denatured salmon sperm DNA and 50°C) and washes (sequentially with 2X SSPE, 0.1% SDS at room temperature for 5 min and then again with fresh solution for 10 min, and finally with 0.5X SSPE, 0.1% SDS at 50°C for 5 min.) were used. Approximately 17 strongly-hybridizing and 17 weakly-hybridizing plaques were identified in the primary screen. Four of the weakly-hybridizing plaques were picked and subjected to one or two further rounds of screening with the radiolabeled probe as above until they were pure. To ensure that these were not delta-15 desaturase clones, they were further analyzed to determine whether they hybridized to an 18 bp oligomer specific to the 3' non-coding region of delta-15 desaturase cDNA (pCF3). After autoradiography of the

SECRET - 5661380

filters, one of the clones was found not to hybridize to this probe. This clone was picked, and a plasmid clone containing the cDNA insert was obtained as described above. Restriction analysis of this plasmid, designated pCM2, showed that it had an approximately 1.3 kb cDNA insert which lacked a 0.7 kb Nco I - Bgl II fragment characteristic of the Arabidopsis delta-15 desaturase cDNA of pCF3. (This fragment corresponds to the DNA located between the Nco I site at nucleotides 474 to 479 and the Bgl II site at nucleotides 1164 to 1169 in SEQ ID NO:1). Partial nucleotide sequences of single strands from the 5' region and 3' region of pCM2 revealed that the cDNA insert was incomplete and that it encoded a polypeptide that is similar to, but distinct from, that encoded by the cDNA in pCF3. In order to isolate a full-length version of the cDNA in plasmid pCM2, the 1.3 kb Not I fragment from plasmid pCM2 containing the cDNA insert was isolated and used as a radiolabeled hybridization probe to rescreen the same Arabidopsis cDNA library as above. Three strongly hybridizing plaques were purified and the plasmids excised as described previously. The three resultant plasmids were digested by Not I restriction enzyme and shown to contain cDNA inserts ranging in size between 1 kB and 1.5 kB. Complete nucleotide sequence determination of the cDNA insert in one of these plasmids, designated pACF2-2, is shown in SEQ ID NO:4. SEQ ID NO:4 shows the 5' to 3' nucleotide sequence of base pairs of the Arabidopsis thaliana cDNA which encodes a fatty acid desaturase. Nucleotides 10-12 and nucleotides 1358 to 1350 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 10 to 1350). The open reading frame was confirmed by comparison of its deduced amino acid sequences with that of the related delta-15

00193 122 566T250

fatty acid desaturase from soybean in this application. Nucleotides 1 to 9 and 1351 to 1525 are, respectively, the 5' and 3' untranslated nucleotides. The 446 amino acid protein sequence in SEQ ID NO:5 is that deduced from the open reading frame in SEQ ID NO:4 and has an estimated molecular weight of 51 kD. Alignment of SEQ ID NOS:2 and 5 shows an overall homology of approximately 80% and that the former has an approximately 55 amino acid long N-terminal extension, which is deduced to be a transit peptide found in nuclear-encoded plastid proteins.

To analyse the developmental expression of the gene corresponding to SEQ ID NO:4, this sequence was used as a radiolabeled hybridization probe on mRNA samples from leaf, root, germinating seedling, and developing siliques from both wild type and mutant line 3707 Arabidopsis plants, essentially as described in Maniatis et al. [Molecular Cloning, A Laboratory Manual (1982) Cold Spring Harbor Laboratory Press]. The results indicated that, in contrast to the constitutive expression of the gene encoding SEQ ID NO:1, the mRNA corresponding to SEQ ID NO:4 is abundant in green tissues, rare in roots and leaves, and is about three-fold more abundant in leaf than that of SEQ ID NO:1. The cDNA in plasmid pCM2 was also shown to hybridize polymorphically to genomic DNA from Arabidopsis thaliana (ecotype Wassileskija and marker line W100 ecotype Landesberg background) digested with Eco RI. It was used as a RFLP marker to map the genetic locus for the gene encoding this fatty acid desaturase in Arabidopsis. A single genetic locus was positioned corresponding to this desaturase cDNA. Its location was thus determined to be on chromosome 3 between the lambda AT228 and cosmid c3838 RFLP markers, "north" of the glabrous locus (Chang et al., Proc. Natl. Acad. Sci. USA (1988))

09249935-12209
SECRET

85:6856-6860; Nam et al., Plant Cell (1989) 1:699-705). This approximates the region to which Arabidopsis fatty acid desaturase fad 2, fad D, and fad B mutations map [Somerville et al., (1992) in press]. Unsuccessful efforts to clone the microsomal delta-12 fatty acid desaturase using cDNA inserts of pCF3 and pACF2-2 alongwith the above data led Applicants to conclude that the cDNA in pACF2-2 encodes a plastid delta-15 fatty acid desaturase that corresponds to the fad D locus.

10 This conclusion will be confirmed by biological expression of the cDNA in pACF2-2.

Plasmid pCM2 was deposited on November 27, 1991 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 68852.

The 1.4 kb, 1.3 kb, and 1.5 kb Not I cDNA insert fragments isolated from plasmids pCF3, pCM2 and pACF2-2 were purified, radiolabeled, and used several times to screen at low stringency as described above two different cDNA libraries: one was made to poly A⁺ mRNA from 3-day-old etiolated Arabidopsis thaliana ("etiolated" library) as described above and one made to polyA⁺ mRNA from the above-ground parts of Arabidopsis thaliana plants, which varied in size from those that had just opened their primary leaves to plants which had bolted and were flowering [Elledge et al. (1991) Proc. Natl. Acad Sci. USA 88:1731-1735]. The cDNA inserts in the library were made into an Xho I site flanked by Eco RI sites in lambda Yes vector [Elledge et al. (1991) Proc. Natl. Acad Sci. USA 88:1731-1735] ("leaf" library). Several plaques from both libraries that hybridized weakly and in duplicate lifts to both SEQ ID NOS:1 and 4 were subjected to plaque purification. Phagemids were excised from the pure phages from "etiolated" library as described above. Plasmids were

356321 356321 356321

5 In all cases, nucleotide sequencing of the cloned DNA revealed clones either identical to SEQ ID NOS:1 or 4 or unrecognizable sequences.

In another set of experiments ca. 400,000 phages in the "leaf" library was screened with SEQ ID NOS:1 and 4 at low stringency (26 C, 1 M Na+, 50% formamide) and high stringency (42 C, 1 M Na+, 50% formamide). Of the several positive signals on the primary plaque lifts, 11 showed high stringency hybridization to SEQ ID NO:1, 35 showed high stringency hybridization to SEQ ID NO:4, and 39 hybridized to both at low stringency only. Twenty seven plaques of the low stringency signals came through a secondary low-stringency screen, 17 of which were used to make DNA from excised plasmids. Of the 7 plasmid DNA were sequenced, 8 were unrecognizable sequences, 5 were identical to SEQ ID NO:1, 2 were identical to SEQ ID NO:2, and 2 were identical to one another and related but distinct to SEQ ID NOS:1 and 4. The novel desaturase sequence, designated pFad-x2, was also isolated from the "leaf" library independently by using as a hybridization probe a 0.6 kB PCR product derived by polymerase chain reaction on poly A+ RNA made from both canola seed as well as Arabidopsis leaves, as described elsewhere in this application, using degenerate oligomers made to conserved sequences between plant delta-15 desaturases and the cyanobacterial des A desaturase. The PCR-derived plasmid, designated pYacp7, was sequenced partially from both ends. Comparison of the sequences of pFad-x2 and pYacp7 revealed that the two independently cloned cDNAs contained an identical sequence that was related to the other delta-15

desaturases and that both were incomplete cDNAs. A partial composite sequence derived from both plasmids, pFadx-2 and pYacp7, is shown in SEQ ID NO:16 as a 5' to 3' nucleotide sequence of 472 bp. Nucleotides 2-4 and nucleotides 468 to 470 are, respectively, the first and the last codons in the open reading frame. This open reading frame is shown in SEQ ID NO:17. Comparison of SEQ ID NO:17 to the other delta-15 desaturase polypeptides disclosed in this application by the method of Needleman et al. [J. Mol. Biol. (1970) 48:443-453] using gap weight and gap length weight values of 3.0 and 0.1, respectively. The overall identities are between 65% and 68% between SEQ ID NO:17 and the microsomal delta-15 desaturases from Arabidopsis, canola and soybean and the overall identities are between 77% and 87% between SEQ ID NO:17 and the plastid delta-15 desaturases from Arabidopsis, canola and soybean. In addition SEQ ID NO:17 has an N-terminal peptide extension compared to the microsomal delta-15 desaturases that shows homology of the transit peptide sequence in Arabidopsis plastid delta-15 desaturase. On the basis of these comparisons it is deduced that SEQ ID NO:16 encodes a plastid delta-15 desaturase. There is genetic data in Arabidopsis suggesting the presence of two loci for plastid delta-15 desaturase. The full-length version of SEQ ID NO:16 can be readily isolated by one skilled in the art. The biological effect of introducing SEQ ID NO:16 or its full-length version into plants will be used to confirm its identity.

30 Plasmid pYacp7 was deposited on 20 November 1992 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 69129.

Using Arabidopsis Delta-15 Desaturase cDNAs
as Hybridization Probes to Isolate
Delta-15 Desaturase cDNAs from Other Plant Species

For the purpose of cloning the Brassica napus seed
5 cDNAs encoding delta-15 fatty acid desaturases, the cDNA
inserts from pCF3 and pCM2 were isolated by polymerase
chain reaction from the respective plasmids,
radiolabeled, and used as hybridization probes to screen
a lambda phage cDNA library made with poly A⁺ mRNA from
10 developing Brassica napus seeds 20-21 days after
pollination. This cDNA library was screened several
times at low stringency, using the Arabidopsis cDNA
probes mentioned above. One of the Brassica napus
cDNAs obtained in the initial screens was used as probe
15 in a subsequent high stringency screen.

Arabidopsis pCM2 insert was radiolabeled and used
as probe to screen approximately 300,000 plaques under
low stringency hybridization conditions. The filter
hybridizations were performed in 50 mM Tris pH 7.6, 6X
20 SSC, 5X Denhardt's, 0.5% SDS, 100 ug denatured calf
thymus DNA at 50°C overnight, and the posthybridization
washes were carried out in 6X SSC, 0.5% SDS at room
temperature for 15 min, then repeated with 2X SSC, 0.5%
SDS at 45°C for 30 min, and then repeated twice with
25 0.2X SSC, 0.5% SDS at 50°C for 30 min. Five strongly-
hybridizing phages were obtained. These were plaque
purified and used to excise the phagemids as described
in the manual of the pBluescriptII Phagemid Kit from
Stratagene (Stratagene 1991 catalogue, item 212205).
30 One of these, designated pBNSF3-2, contained a 1.3 kb
insert. pBNSF3-f2 was sequenced completely on both
strands and the nucleotide sequence is shown in SEQ ID
NO:6. Plasmid pBNSF3-2 was deposited on 27 November
1991 with the American Type Culture Collection of

Rockville Maryland, USA under the provisions of the Budapest Treaty and bears the accession number 68854.

An additional low stringency screen using pCM2 probe provided eight strongly hybridizing phages. One of these, designated pBNSFd 8, contained a 0.4kb insert. pBNSFd-8 was sequenced completely on one strand, this nucleotide sequence showed significant divergence from the sequence SEQ ID NO:6 in the homologous region, which suggested that it corresponded to a novel Brassica napus seed desaturase different from that shown in SEQ ID NO:6. pBNSFd-8 insert was radiolabelled and used as hybridization probe in a high stringency screen of the Brassica napus seed cDNA library. The hybridization conditions were identical to those of the low stringency screen described above except for the temperature of the final two 30 min posthybridization washes in 0.2x SSC, 0.5% SDS was increased to 60°C. This screen resulted in three strongly hybridizing phages that were purified and excised. One of the excised plasmids pBNSFd-3 contained a 1.4kb insert that was sequenced completely on both strands. SEQ ID NO:8 shows the complete nucleotide sequence of pBNSFd-2.

Using Arabidopsis Delta-15 Desaturase cDNA as a Hybridization Probe to Isolate a Glycerolipid

Desaturase cDNA from Soybean

A cDNA library was made to poly A⁺ mRNA isolated from developing soybean seeds, and screened essentially as described above, except that filters were prehybridized in 25 mL of hybridization buffer consisting of 50mM Tris-HCl, pH 7.5, 1 M NaCl, 1% SDS, 5% dextran sulfate and 0.1 mg/mL denatured salmon sperm DNA (Sigma Chemical Co.) at 50°C for 2 h. Radiolabeled probe prepared from pCF3 as described above was added, and allowed to hybridize for 18 h at 50°C. The probes were washed twice at room temperature with 2X SSPE, 1%

862221 5E6F260

SDS for five min followed by washing for 5 min at 50°C in 0.2X SSPE, 1% SDS. Autoradiography of the filters indicated that there was one strongly hybridizing plaque, and approximately five weakly hybridizing
5 plaques. The more strongly hybridizing plaque was subjected to a second round of screening as before, except that the final wash was for 5 min at 60°C in 0.2X SSPE, 1% SDS. Numerous, strongly hybridizing plaques were observed, and one, well-isolated from other phage,
10 was picked for further analysis.

Sequences of the pBluescript vector from the purified phage, including the cDNA insert, were excised in the presence of a helper phage and the resultant phagemid was used to infect *E. coli* XL-1 Blue cells.
15 DNA from the plasmid, designated pXF1, was made by the alkaline lysis miniprep procedure described in Sambrook et al. (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press). The alkali-denatured double-stranded DNA from pXF1 was
20 completely sequenced on both strands. The insert of pXF1 contained a stretch of 1783 nucleotides which contained an unknown open-reading frame and also contained a poly-A stretch of 16 nucleotides 3' to the open reading frame, from nucleotides 1767 to 1783,
25 followed by an Eco RI restriction site. The 2184 bases that followed this Eco RI site contained a 1145 bp open reading frame which encoded a polypeptide of about 68% identity to, and colinear with, the Arabidopsis delta-15 desaturase polypeptide listed in SEQ ID No:2. The
30 putative start methionine of the 1145 bp open-reading frame corresponded to the start methionine of the Arabidopsis microsomal delta-15 peptide and there were no amino acids corresponding to a plastid transit peptide 5' to this methionine. When the insert in pXF1
35 was digested with Eco RI four fragments were observed,

fragments of approximately 370 bp and 1400 bp fragments,
derived from the first 1783 bp of the insert in pXF1,
and fragments of approximately 600 bp and 1600 bp
derived from the the other 2184 nucleotides of the
5 insert in pXF1. Only the 600 bp and 1600 bp fragments
hybridized with probe derived from pCF3 on Southern
blots. It was deduced that pXF1 contained two different
cDNA inserts separated by an Eco RI site and the second
of these inserts was a 2184 bp cDNA encoding a soybean
10 microsomal delta-15 desaturase. The complete nucleotide
sequence of the 2184 bp soybean microsomal delta-15 cDNA
contained in plasmid pXF1 is listed in SEQ ID No:10.
Plasmid pXF1 was deposited on December 3, 1991 with the
American Type Culture Collection of Rockville, Maryland
15 under the provisions of the Budapest Treaty and bears
accession number ATCC 68874.

Using Soybean Microsomal Delta-15 Desaturase cDNA as a
Hybridization Probe to Isolate
cDNAs Encoding Related Desaturases from Soybean

20 A 1.0 kb fragment of DNA corresponding to part of
the coding region of the soybean microsomal delta-15
desaturase cDNA contained in plasmid pXF1, was excised
with the restriction enzyme Hha I and gel purified. The
fragment was labeled with ³²P as described above and
25 used to probe a soybean cDNA library as described above.
Autoradiography of the filters indicated that there were
eight hybridizing plaques and these were subjected to a
second round of screening. Sequences of the pBluescript
vector from all eight of the purified phages, including
30 the cDNA inserts, were excised in the presence of a
helper phage and the resultant phagemids were used to
infect *E. coli* XL-1 Blue cells. DNA from the plasmids
was made by the alkaline lysis miniprep procedure
described in Sambrook et al. (Molecular Cloning, A
35 Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor

000193313300
000000000000

Laboratory Press). Restriction analysis showed they contained inserts ranging from 1.0 kb to 3.0 kb in size. One of these inserts, designated pSFD-118bwp, contained an insert of about 1700 bp. The alkali-denatured
5 double-stranded DNA from pSFD-118bwp was completely sequenced on both strands, shown in SEQ ID NO:12. The insert of pSFD-118bwp contained a stretch of 1675 nucleotides which contained an open-reading frame encoding a polypeptide, shown in SEQ ID NO:13, of about
10 80% identity with, and colinear with, the Arabidopsis plastid delta-15 desaturase polypeptide listed in SEQ ID No:5. The open-reading frame also encoded amino acids corresponding to a plastid transit peptide at the 5' end of the open-reading frame. The transit peptide was
15 colinear with, and shared some homology to, the transit peptide described for the Arabidopsis plastid delta-15 glycerolipid desaturase. The complete nucleotide sequence of the 1675 bp soybean plastid delta-15 glycerolipid desaturase cDNA is listed in SEQ ID No:12.
20 Comparison of the different delta-15 desaturase sequences disclosed in the application by the method of Needleman et al. (J. Mol. Biol. (1970) 48:443-453) using gap weight and gap length weight values of 3.0 and 0.1, respectively, reveals the relatedness between them as shown
25 in Table 3.

SECRET 3667260

TABLE 3

Percent Identities Between Different Delta-15
Fatty Acid Desaturases at the Amino Acid Level

	<u>aD</u>	<u>c3</u>	<u>cD</u>	<u>s3</u>	<u>sD</u>
a3	66	93	66	68	67
aD	-	67	90	67	69
c3	-	-	68	68	68
cD	-	-	-	68	74

a3, aD, c3, cD, s3 and sD refer, respectively, to
SEQ ID NO:2 (Arabidopsis microsomal delta-15
desaturase), SEQ ID NO:5 (Arabidopsis plastid delta-15
desaturase), SEQ ID NO:7 (canola microsomal delta-15
desaturase), SEQ ID NO:9 (canola plastid delta-15
desaturase), SEQ ID NO:11 (soybean microsomal delta-15
desaturase), and SEQ ID NO:13 (soybean plastid delta-15
desaturase). Based on these comparisons, the delta-15
desaturases, of both microsomal and plastid types, have
overall identities of 65% or more at the amino acid
levels, even when from different plant species.

Isolation of Nucleotide Sequences Encoding
Homologous and Heterologous Glycerolipid Desaturases

Fragments of the instant invention may be used to
isolate cDNAs and genes of homologous and heterologous
glycerolipid desaturases from the same species as the
fragment of the invention or from different species.
Isolation of homologous genes using sequence-dependent
protocols is well-known in the art. Southern blot
analysis revealed that the Arabidopsis microsomal
delta-15 desaturase cDNA (SEQ ID NO:1) hybridized to
genomic DNA fragments of corn and soybean. In addition,
Applicants have demonstrated that it can be used to
isolate cDNAs encoding seed microsomal delta-15
desaturases from Brassica napus (SEQ ID NO:6) and
soybean (SEQ ID NO:10). Thus, one can isolate cDNAs and

SECRET "SECRET"

genes for homologous glycerolipid desaturases from the same or different higher plant species, especially from the oil-producing species.

More importantly, one can use the fragments of the invention to isolate cDNAs and genes for heterologous glycerolipid desaturases, including those found in plastids. Thus, Arabidopsis microsomal delta-15 desaturase cDNA (SEQ ID NO:1) was successfully used as a hybridization probe to isolate cDNAs encoding the related plastid delta-15 desaturases from Arabidopsis (SEQ ID NO:4) and Brassica napus (SEQ ID NO: 8), and the soybean microsomal delta-15 soybean (SEQ ID NO:10) was successfully used to isolate soybean cDNA encoding plastid delta-15 desaturase (SEQ ID NO:12).

In a particular embodiment of the present invention, regions of the nucleic acid fragments of the invention that are conserved between different desaturases may be used by one skilled in the art to design a mixture of degenerate oligomers for use in sequence-dependent protocols aimed at isolating nucleic acid fragments encoding other homologous or heterologous glycerolipid desaturase cDNA's or genes. For example, by comparing all desaturase polypeptides one can identify stretches of amino acids that are conserved between them, and then use the conserved amino acid sequence to design oligomers, both short degenerate or long ones, or "guessmers" as known by one skilled in the art (see Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press). Such oligomers and "guessmers" may be used as hybridization probes as known to one skilled in the art.

For example, comparison of cyanobacterial desA and plant delta-15 desaturases revealed a particularly well conserved stretch of amino acids (amino acids 97-108 in

SEQ ID NO:1). SEQ ID NOS:20 and 21 represent two sets of 36-mers each 16-fold degenerate made to this region. End-labeled oligomers represented in SEQ ID NOS:20 and 21 were mixed and used as hybridization probes to screen Arabidopsis cDNA libraries. Most of the positively-hybridizing plaques also hybridized to cDNAs encoding Arabidopsis microsomal and plastid delta-15 desaturases (SEQ ID NOS:1 and 4). However, the use of SEQ ID NOS:20 and 21 did not give consistent and reproducible results. A 135 base-long oligomer (SEQ ID NO:32) was also made as an antisense strand to a longer stretch of the same conserved region, amino acids 97 to 141 in SEQ ID NO:1 (FVLGHDCGHGSFSDIPLLNSVVGHILHSFILVPYHGWRISHRTHH). At positions of ambiguity, the design used either deoxyinosines or most frequently used codons based on the codon usage in Arabidopsis genes. When used as a hybridization probe, the 135-mer hybridized to all plaques that also hybridized to cDNAs encoding Arabidopsis microsomal and plastid delta-15 desaturases (SEQ ID NOS:1 and 4). In addition, it also hybridized to plaques that did not hybridize to SEQ ID NOS:1 and 4). The latter were purified and excised as described previously. Nucleotide sequencing of the cDNA inserts in the resultant plasmids revealed DNA sequences that did not show any relatedness to any desaturase.

For another example, in the polymerase chain reaction (Innis, et al., Eds, (1990) PCR Protocols: A Guide to Methods and Applications, Academic Press, San Diego), two short pieces of the present fragment of the invention can be used to amplify a longer glycerolipid desaturase DNA fragment from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleotide sequences with one primer based on the fragment of the invention and the other on either the poly A⁺ tail or a vector sequence. These oligomers may

86221-5667208

be unique sequences or degenerate sequences derived from the nucleic acid fragments of the invention. The longer piece of homologous glycerolipid desaturase DNA generated by this method could then be used as a probe

5 for isolating related glycerolipid desaturase genes or cDNAs from Arabidopsis or other species. The design of oligomers, including long oligomers using deoxyinosine, and "guessmers" for hybridization or for the polymerase chain reaction are known to one skilled in the art and

10 discussed in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press). Stretches of conserved amino acids between delta-15 desaturase and other desaturases, especially desA, allow for the design of such oligomers.

15 For example, conserved stretches of amino acids between desA and delta-15 desaturase, discussed above, are useful in designing long oligomers for hybridization as well as shorter ones for use as primers in the polymerase chain reaction. In this regard, the

20 conserved amino acid stretch of amino acids 97 to 108 of SEQ ID NO:2 is particularly useful. Other conserved regions in SEQ ID NO:2 useful for this purpose are amino acids 299 to 309, amino acids 115 to 121, and amino acids 133 to 141. Amino acid stretch 133 to 141 in SEQ

25 ID NO:2 shows especially good homology to several desaturases. For example, in this stretch, amino acids 133, 137, 138, 140 and 141 are conserved in plant delta-15 desaturases, cyanobacterial desA, yeast and mammalian microsomal stearyl-CoA desaturases.

30 Comparison of cyanobacterial des A and plant delta-15 desaturases revealed two particularly well conserved stretch of amino acids (amino acids 97-108 and amino acids 299-311 in SEQ ID NO:1) that can be used for PCR. The following sets of PCR primers were made to these

35 regions:

SEQ ID NO	Length	Fold Degeneracy	AA positions in SEQ ID NO:2	AA Sequence
20	36	16	97-108 (S)	FVLGHDCGHGSF
21	36	16	97-108 (S)	FVLGHDCGHGSF
28	36	16	97-108 (S)	FVLGHDCGHGSF
29	36	16	97-108 (S)	FVLGHDCGHGSF
22	18	72	100-105 (S)	GHDCGH
23	18	72	100-105 (S)	GHDCGH
24	18	72	299-304 (AS)	HDIGTH
25	18	72	299-304 (AS)	HDIGTH
26	23	416	304-309 (AS)	HVIHHL
27	23	416	304-309 (AS)	HVIHHL
30	38	64	299-311 (AS)	HDIGTHVHHHLFP
31	38	64	299-311 (AS)	HDIGTHVHHHLFP

- In one experiment, PCRs were performed using SEQ ID NOS:22 and 23 as sense primers and either SEQ ID NOS:24 and 25 or SEQ ID NOS:26 and 27 as antisense primers on poly A+ RNA purified from both Arabidopsis leaf and canola developing seeds. All PCRs resulted in PCR products of the correct size (ca. 630 bp). The PCR products from Arabidopsis and canola were purified and used as radiolabeled hybridization probes to screen the Lambda Yes Arabidopsis cDNA library, as described above.
- 5 This led to the isolation of a pure phage, which was excised to give plasmid pYacp7. The cDNA insert in pYacp7 was partially sequenced. It's sequence showed that it encoded an incomplete desaturase polypeptide that was identical to another cDNA (in plasmid pFadx-2)
- 10 isolated by low-stringency hybridization as described previously. The composite sequence derived from the partial sequences from the cDNA inserts in pFadx-2 and pYacp7 is shown in SEQ ID NO:16 and the polypeptide encoded by it in SEQ ID NO:17. As discussed previously,
- 15 SEQ ID NO:17 is a putative plastid delta-15 desaturase. This is further supported by Southern blot analysis
- 20

using radiolabeled cDNA inserts from either pCF3, pACF2-2, or pYacp7 on Arabidopsis genomic DNA digested with one of several enzymes. It shows that the different inserts hybridize to different restriction fragments and that only the inserts from pACF2-2 and pYacp7 show some cross-hybridization.

In another PCR experiment, PCR was performed using ca. 80 pmoles each of SEQ ID NOS:28 and 29 as sense primers and ca. 94 pmoles each of SEQ ID NOS:30 and 31 as antisense primers on poly A+ RNA purified from Arabidopsis mutant line 3707. This was performed using GeneAmp® RNA PCR Kit (Perkin Elmer Cetus) following manufacturer's protocol and using the following program: a) 1 cycle of 2 min at 95°C, b) 35 cycles of 1 min at 95°C (denaturation), 1 min at 50°C (annealing) and 1 min at 65°C (extension), and c) 1 cycle of 7 min at 65°C. The resulting PCR product, of the correct size (ca. 630 bp), was purified, radiolabeled, and used as a hybridization probe on a Southern blot of Arabidopsis genomic DNA as described above. While it hybridized to restriction fragments that also hybridized to SEQ ID NOS:1 (Arabidopsis microsomal delta-15 desaturase), 4 (Arabidopsis plastid delta-15 desaturase), and 16 (Arabidopsis plastid delta-15 desaturase), it also hybridized to novel fragments that did not hybridize to previously cloned desaturase cDNAs. However, even after several attempts, the radiolabeled PCR product did not hybridize to any novel cDNA clone when used as a probe on different Arabidopsis cDNA libraries: in all cases it hybridized only to plaques that also hybridized to the known desaturase cDNAs. Furthermore, the PCR product was subcloned into a plasmid vector and after screening about a 100 of these, none gave rise to a clone with a novel desaturase sequence.

25221 5554250
The isolation of other glycerolipid desaturases will become easier as more examples of glycerolipid desaturases are isolated using the fragments of the invention. Knowing the conserved amino acid sequences from diverse desaturases will also allow one to identify more and better consensus sequences. Such sequences can be used to make hybridization probes or amplification primers which will further aid in the isolation of different glycerolipid desaturases, including those from non-plant sources such as fungi, algae, and even cyanobacteria, as well as other membrane-associated desaturases from other organisms.

The function of the diverse nucleotide fragments encoding glycerolipid desaturases that can be isolated using the present invention can be identified by transforming plants with the isolated desaturase sequences, linked in sense or antisense orientation to suitable regulatory sequences required for plant expression, and observing the fatty acid phenotype of the resulting transgenic plants. Preferred target plants for the transformation are the same as the source of the isolated nucleotide fragments when the goal is to obtain inhibition of the corresponding endogenous gene by antisense inhibition or cosuppression. Preferred target plants for use in expression or overexpression of the isolated nucleic acid fragments are plants with known mutations in desaturation reactions, such as the Arabidopsis desaturase mutants, mutant flax deficient in delta-15 desaturation, or mutant sunflower deficient in delta-12 desaturation. Alternatively, the function of the isolated nucleic acid fragments can be determined similarly via transformation of other organisms, such as yeast or cyanobacteria, with chimeric genes containing the nucleic acid fragment and suitable regulatory

sequences followed by analysis of fatty acid composition and/or enzyme activity.

Overexpression of the Glycerolipid
Desaturase Enzymes in Transgenic Species

5 The nucleic acid fragment(s) of the instant invention encoding functional glycerolipid desaturase(s), with suitable regulatory sequences, can be used to overexpress the enzyme(s) in transgenic organisms. Such recombinant DNA constructs may include
10 either the native glycerolipid desaturase gene or a chimeric glycerolipid desaturase gene isolated from the same or a different species as the host organism. For overexpression of glycerolipid desaturase(s), it is preferable that the introduced gene be from a different
15 species to reduce the likelihood of cosuppression. For example, overexpression of delta-15 desaturase in soybean, rapeseed, or other oil-producing species to produce altered levels of polyunsaturated fatty acids may be achieved by expressing RNA from the entire cDNA
20 found in pCF3. Similarly, the isolated nucleic acid fragments encoding glycerolipid desaturases from Arabidopsis, rapeseed, and soybean can also be used by one skilled in the art to obtain substantially homologous full-length cDNAs, if not already obtained,
25 as well as the corresponding genes as fragments of the invention. These, in turn, may be used to overexpress the corresponding desaturases in plants. One skilled in the art can also isolate the coding sequence(s) from the fragment(s) of the invention by using and/or creating
30 sites for restriction endonucleases, as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press). For example, the fragment in SEQ ID NO:1 in plasmid pCF3 is flanked by Not I sites and can be
35 isolated as a Not I fragment that can be introduced in

the sense orientation relative to suitable plant regulatory sequences. Alternatively, sites for Nco I (5'-CCATGG-3') or Sph I (5'-GCATGC-3') that allow precise removal of coding sequences starting with the
 5 initiating codon "ATG" may be engineered into the fragment(s) of the invention. For example, for utilizing the coding sequence of delta-15 desaturase from pCF3, an Sph I site can be engineered by substituting nucleotides at positions 44, 45, and 49 of
 10 SEQ ID NO:1 with G, C, and C, respectively.

Inhibition of Plant Target Genes by Use of Antisense RNA

Antisense RNA has been used to inhibit plant target genes in a tissue-specific manner (see van der Krol et
 15 al., Biotechniques (1988) 6:958-976). Antisense inhibition has been shown using the entire cDNA sequence (Sheehy et al., Proc. Natl. Acad. Sci. USA (1988) 85:8805-8809) as well as a partial cDNA sequence (Cannon et al., Plant Molec. Biol. (1990) 15:39-47). There is
 20 also evidence that the 3' non-coding sequences (Ch'ng et al., Proc. Natl. Acad. Sci. USA (1989) 86:10006-10010) and fragments of 5' coding sequence, containing as few as 41 base-pairs of a 1.87 kb cDNA (Cannon et al., Plant Molec. Biol. (1990) 15:39-47), can
 25 play important roles in antisense inhibition.

The use of antisense inhibition of the glycerolipid desaturases may require isolation of the transcribed sequence for one or more target glycerolipid desaturase genes that are expressed in the target tissue of the
 30 target plant. The genes that are most highly expressed are the best targets for antisense inhibition. These genes may be identified by determining their levels of transcription by techniques, such as quantitative analysis of mRNA levels or nuclear run-off
 35 transcription, known to one skilled in the art.

001995-1333

For example, antisense inhibition of delta-15 desaturase in Brassica napus resulting in altered levels of polyunsaturated fatty acids may be achieved by expressing antisense RNA from the entire or partial cDNA found in pBNSF3-2.

Inhibition of Plant Target Genes by Cosuppression

The phenomenon of cosuppression has also been used to inhibit plant target genes in a tissue-specific manner. Cosuppression of an endogenous gene using the entire cDNA sequence (Napoli et al., The Plant Cell (1990) 2:279-289; van der Krol et al., The Plant Cell (1990) 2:291-299) as well as a partial cDNA sequence (730 bp of a 1770 bp cDNA) (Smith et al., Mol. Gen. Genetics (1990) 224:477-481) are known.

The nucleic acid fragments of the instant invention encoding glycerolipid desaturases, or parts thereof, with suitable regulatory sequences, can be used to reduce the level of glycerolipid desaturases, thereby altering fatty acid composition, in transgenic plants which contain an endogenous gene substantially homologous to the introduced nucleic acid fragment. The experimental procedures necessary for this are similar to those described above for the overexpression of the glycerolipid desaturase nucleic acid fragments. For example, cosuppression of delta-15 desaturase in Brassica napus resulting in altered levels of polyunsaturated fatty acids may be achieved by expressing in the sense orientation the entire or partial seed delta-15 desaturase cDNA found in pBNSF3-2.

Selection of Hosts, Promoters and Enhancers

A preferred class of heterologous hosts for the expression of the nucleic acid fragments of the invention are eukaryotic hosts, particularly the cells of higher plants. Particularly preferred among the

higher plants are the oil-producing species, such as soybean (Glycine max), rapeseed (including Brassica napus, B. campestris), sunflower (Helianthus annuus), cotton (Gossypium hirsutum), corn (Zea mays), cocoa
 5 (Theobroma cacao), safflower (Carthamus tinctorius), oil palm (Elaeis guineensis), coconut palm (Cocos nucifera), flax (Linum usitatissimum), and peanut (Arachis hypogaea).

Expression in plants will use regulatory sequences
 10 functional in such plants. The expression of foreign genes in plants is well-established (De Blaere et al., Meth. Enzymol. (1987) 153:277-291). The source of the promoter chosen to drive the expression of the fragments of the invention is not critical provided it has
 15 sufficient transcriptional activity to accomplish the invention by increasing or decreasing, respectively, the level of translatable mRNA for the glycerolipid desaturases in the desired host tissue. Preferred promoters include (a) strong constitutive plant
 20 promoters, such as those directing the 19S and 35S transcripts in cauliflower mosaic virus (Odell et al., Nature (1985) 313:810-812; Hull et al., Virology (1987) 86:482-493), and (b) tissue- or developmentally-specific promoters. Examples of tissue-specific promoters are
 25 the light-inducible promoter of the small subunit of ribulose 1,5-bis-phosphate carboxylase (if expression is desired in photosynthetic tissues), the maize zein protein promoter (Matzke et al., EMBO J. (1984) 3:1525-1532), and the chlorophyll a/B binding protein
 30 promoter (Lampa et al., Nature (1986) 316:750-752).

Particularly preferred promoters are those that allow seed-specific expression. This may be especially useful since seeds are the primary source of vegetable oils and also since seed-specific expression will avoid
 35 any potential deleterious effect in non-seed tissues.

SECRET 5667289

Examples of seed-specific promoters include, but are not limited to, the promoters of seed storage proteins, which can represent up to 90% of total seed protein in many plants. The seed storage proteins are strictly regulated, being expressed almost exclusively in seeds in a highly tissue-specific and stage-specific manner (Higgins et al., Ann. Rev. Plant Physiol. (1984) 35:191-221; Goldberg et al., Cell (1989) 56:149-160). Moreover, different seed storage proteins may be expressed at different stages of seed development.

Expression of seed-specific genes has been studied in great detail (See reviews by Goldberg et al., Cell (1989) 56:149-160 and Higgins et al., Ann. Rev. Plant Physiol. (1984) 35:191-221). There are currently numerous examples of seed-specific expression of seed storage protein genes in transgenic dicotyledonous plants. These include genes from dicotyledonous plants for bean b-phaseolin (Sengupta-Gopalan et al., Proc. Natl. Acad. Sci. USA (1985) 82:3320-3324; Hoffman et al., Plant Mol. Biol. (1988) 11:717-729), bean lectin (Voelker et al., EMBO J. (1987) 6:3571-3577), soybean lectin (Okamuro et al., Proc. Natl. Acad. Sci. USA (1986) 83:8240-8244), soybean Kunitz trypsin inhibitor (Perez-Grau et al., Plant Cell (1989) 1:095-1109), soybean b-conglycinin (Beachy et al., EMBO J. (1985) 4:3047-3053; pea vicilin (Higgins et al., Plant Mol. Biol. (1988) 11:683-695), pea convicilin (Newbiggin et al., Planta (1990) 180:461-470), pea legumin (Shirsat et al., Mol. Gen. Genetics (1989) 215:326-331); rapeseed napin (Radke et al., Theor. Appl. Genet. (1988) 75:685-694) as well as genes from monocotyledonous plants such as for maize 15 kD zein (Hoffman et al., EMBO J. (1987) 6:3213-3221), maize 18 kD oleosin (Lee et al., Proc. Natl. Acad. Sci. USA (1991) 888:6181-6185), barley b-hordein (Marris et al., Plant Mol. Biol. (1988)

3522T SE6460

5 their temporal and spatial expression pattern in
transgenic plants. Such examples include use of
Arabidopsis thaliana 2S seed storage protein gene
promoter to express enkephalin peptides in Arabidopsis
and B. napus seeds (Vandekerckhove et al.,
10 Bio/Technology (1989) 7:929-932), bean lectin and bean
b-phaseolin promoters to express luciferase (Riggs et
al., Plant Sci. (1989) 63:47-57), and wheat glutenin
promoters to express chloramphenicol acetyl transferase
(Colot et al., EMBO J. (1987) 6:3559-3564).

15 Of particular use in the expression of the nucleic
acid fragment of the invention will be the heterologous
promoters from several soybean seed storage protein
genes such as those for the Kunitz trypsin inhibitor
(Jofuku et al., Plant Cell (1989) 1:1079-1093; glycinin
20 (Nielson et al., Plant Cell (1989) 1:313-328), and
b-conglycinin (Harada et al., Plant Cell (1989)
1:415-425). Promoters of genes for a- and b-subunits of
soybean β -conglycinin storage protein will be
particularly useful in expressing the mRNA or the
25 antisense RNA in the cotyledons at mid- to late-stages
of seed development (Beachy et al., EMBO J. (1985)
4:3047-3053) in transgenic plants. This is because
there is very little position effect on their expression
in transgenic seeds, and the two promoters show
30 different temporal regulation. The promoter for the
a-subunit gene is expressed a few days before that for
the b-subunit gene. This is important for transforming
rapeseed where oil biosynthesis begins about a week
before seed storage protein synthesis (Murphy et al., J.
35 Plant Physiol. (1989) 135:63-69).

Also of particular use will be promoters of genes expressed during early embryogenesis and oil biosynthesis. The native regulatory sequences, including the native promoters, of the glycerolipid desaturase genes expressing the nucleic acid fragments of the invention can be used following their isolation by those skilled in the art. Heterologous promoters from other genes involved in seed oil biosynthesis, such as those for B. napus isocitrate lyase and malate synthase (Comai et al., Plant Cell (1989) 1:293-300), delta-9 desaturase from safflower (Thompson et al. Proc. Natl. Acad. Sci. USA (1991) 88:2578-2582) and castor (Shanklin et al., Proc. Natl. Acad. Sci. USA (1991) 88:2510-2514), acyl carrier protein (ACP) from Arabidopsis (Post-Beittenmiller et al., Nucl. Acids Res. (1989) 17:1777), B. napus (Safford et al., Eur. J. Biochem. (1988) 174:287-295), and B. campestris (Rose et al., Nucl. Acids Res. (1987) 15:7197), b-ketoacyl-ACP synthetase from barley (Siggaard-Andersen et al., Proc. Natl. Acad. Sci. USA (1991) 88:4114-4118), and oleosin from Zea mays (Lee et al., Proc. Natl. Acad. Sci. USA (1991) 88:6181-6185), soybean (Genbank Accession No: X60773) and B. napus (Lee et al., Plant Physiol. (1991) 96:1395-1397) will be of use. If the sequence of the corresponding genes is not disclosed or their promoter region is not identified, one skilled in the art can use the published sequence to isolate the corresponding gene and a fragment thereof containing the promoter. The partial protein sequences for the relatively-abundant enoyl-ACP reductase and acetyl-CoA carboxylase are also published (Slabas et al.; Biochim. Biophys. Acta (1987) 877:271-280; Cottingham et al., Biochim. Biophys. Acta (1988) 954:201-207) and one skilled in the art can use these sequences to isolate the corresponding seed genes with their promoters. Similarly, the fragments of the

852221 51667260

present invention encoding glycerolipid desaturases can be used to obtain promoter regions of the corresponding genes for use in expressing chimeric genes.

Attaining the proper level of expression of the
5 nucleic acid fragments of the invention may require the use of different chimeric genes utilizing different promoters. Such chimeric genes can be transferred into host plants either together in a single expression vector or sequentially using more than one vector.

10 It is envisioned that the introduction of enhancers or enhancer-like elements into the promoter regions of either the native or chimeric nucleic acid fragments of the invention will result in increased expression to accomplish the invention. This would include viral
15 enhancers such as that found in the 35S promoter (Odell et al., Plant Mol. Biol. (1988) 10:263-272), enhancers from the opine genes (Fromm et al., Plant Cell (1989) 1:977-984), or enhancers from any other source that result in increased transcription when placed into a
20 promoter operably linked to the nucleic acid fragment of the invention.

Of particular importance is the DNA sequence element isolated from the gene for the α -subunit of b-conglycinin that can confer 40-fold seed-specific
25 enhancement to a constitutive promoter (Chen et al., Dev. Genet. (1989) 10:112-122). One skilled in the art can readily isolate this element and insert it within the promoter region of any gene in order to obtain seed-specific enhanced expression with the promoter in
30 transgenic plants. Insertion of such an element in any seed-specific gene that is expressed at different times than the b-conglycinin gene will result in expression in transgenic plants for a longer period during seed development.

09219936-22396

The invention can also be accomplished by a variety of other methods to obtain the desired end. In one form, the invention is based on modifying plants to produce increased levels of glycerolipid desaturases by virtue of introducing more than one copy of the foreign gene containing the nucleic acid fragments of the invention. In some cases, the desired level of polyunsaturated fatty acids may require introduction of foreign genes for more than one kind of glycerolipid desaturase.

Any 3' non-coding region capable of providing a polyadenylation signal and other regulatory sequences that may be required for the proper expression of the nucleic acid fragments of the invention can be used to accomplish the invention. This would include 3' ends of the native glycerolipid desaturase(s), viral genes such as from the 35S or the 19S cauliflower mosaic virus transcripts, from the opine synthesis genes, ribulose 1,5-bisphosphate carboxylase, or chlorophyll a/b binding protein. There are numerous examples in the art that teach the usefulness of different 3' non-coding regions.

Transformation Methods

Various methods of transforming cells of higher plants according to the present invention are available to those skilled in the art (see EPO Pub. 0 295 959 A2 and 0 318 341 A1). Such methods include those based on transformation vectors utilizing the Ti and Ri plasmids of Agrobacterium spp. It is particularly preferred to use the binary type of these vectors. Ti-derived vectors transform a wide variety of higher plants, including monocotyledonous and dicotyledonous plants (Sukhapinda et al., Plant Mol. Biol. (1987) 8:209-216; Potrykus, Mol. Gen. Genet. (1985) 199:183). Other transformation methods are available to those skilled in the art, such as direct uptake of foreign DNA constructs

(see EPO Pub. 0 295 959 A2), techniques of electroporation (Fromm et al., Nature (1986) (London) 319:791) or high-velocity ballistic bombardment with metal particles coated with the nucleic acid constructs (Kline et al., Nature (1987) (London) 327:70). Once transformed, the cells can be regenerated by those skilled in the art.

Of particular relevance are the recently described methods to transform foreign genes into commercially important crops, such as rapeseed (De Block et al., Plant Physiol. (1989) 91:694-701), sunflower (Everett et al., Bio/Technology (1987) 5:1201), and soybean (Christou et al., Proc. Natl. Acad. Sci USA (1989) 86:7500-7504).

15 Application to RFLP Technology

The use of restriction fragment length polymorphism (RFLP) markers in plant breeding has been well-documented in the art (Tanksley et al., Bio/Technology (1989) 7:257-264). The nucleic acid fragments of the invention can be used as RFLP markers for traits linked to expression of glycerolipid desaturases. These traits will include altered levels of unsaturated fatty acids. The nucleic acid fragment of the invention can also be used to isolate the glycerolipid desaturase gene from variant (including mutant) plants with altered levels of unsaturated fatty acids. Sequencing of these genes will reveal nucleotide differences from the normal gene that cause the variation. Short oligonucleotides designed around these differences may be used as hybridization probes to follow the variation in polyunsaturates. Oligonucleotides based on differences that are linked to the variation may be used as molecular markers in breeding these variant oil traits.

EXAMPLES

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. All publications, including patents and non-patent literature, referred to in this specification are expressly incorporated by reference herein.

EXAMPLE 1

ISOLATION OF GENOMIC DNA FLANKING THE T-DNA SITE OF
INSERTION IN ARABIDOPSIS THALIANA MUTANT LINE 3707

20 Identification of an Arabidopsis thaliana T-DNA Mutant
with Low Linolenic Acid Content

A population of Arabidopsis thaliana (geographic race Wassilewskija) transformants containing the T-DNA of Agrobacterium tumefaciens was generated by seed transformation as described by Feldmann et al., (Mol. Gen. Genetics (1987) 208:1-9). In this population the transformants contain DNA sequences encoding the pBR322 bacterial vector, nopaline synthase, neomycin phosphotransferase (NPTII, confers kanamycin resistance), and b-lactamase (confers ampicillin resistance) within the T-DNA border sequences. The integration of the T-DNA into different areas of the chromosomes of individual transformants may cause a disruption of plant gene function at or near the site of insertion, and phenotypes associated with this loss of

gene function can be analyzed by screening the population for the phenotype.

T3 seed was generated from the wild type seed treated with Agrobacterium tumefaciens by two rounds of self-fertilization as described by Feldmann et al., (Science (1989) 243:1351-1354). These progeny were segregating for the T-DNA insertion, and thus for any mutation resulting from the insertion. Approximately 100 seeds of each of 6000 lines were combined and the fatty acid content of each of the 6000 pooled samples was determined by gas chromatography of the fatty acyl methyl esters essentially as described by Browse et al., (Anal. Biochem. (1986) 152:141-145) except that 2.5% H₂SO₄ in methanol was used as the methylation reagent and samples were heated for 1.5 h at 80°C to effect the methanolysis of the seed triglycerides. A line designated "3707" produced seeds that gave an altered fatty acid profile compared to that of the total population. T3 plants were grown from individual T3 seeds produced by line 3707 and self-fertilized to produce T4 seeds on individual plants that were either homozygous wild type, homozygous mutant, or heterozygous for the mutation. The percent fatty acid compositions of a representative subsample of the entire population, of the pooled 3707 T3 seeds, and of a homozygous T4 mutant segregant are shown in Table 4.

TABLE 4

<u>Fatty Acid Methyl Ester</u>	<u>T3 Pools from lines 3501-4000 average and (std. deviation)</u>	<u>3707 T3 Pool</u>	<u>3707 Homozygous T4 Segregant</u>
palmitic	7.4 (0.37)	7.0	6.4
stearic	3.0 (0.22)	2.9	3.0
oleic	17.0 (1.5)	17.7	15.9
linoleic	29.3 (0.78)	35.0	42.4
linolenic	16.1 (1.1)	10.2	3.1
eicosenoic	20.2 (0.73)	20.5	23.6

The phenotype of the segregating T3 pool of line 3707 (high linoleic acid, low linolenic acid) was

intermediate between that of the population subsample

5 and the homozygous T4 mutant seeds suggesting that line 3707 harbored a mutation at a locus which controls the conversion of linoleic to linolenic acid in the seed. Still, it was not apparent whether the mutant phenotype in line 3707 was the result of a T-DNA insertion.

10 Therefore, Applicants checked a segregating T4 population to determine whether the mutant fatty acid phenotype cosegregated with the nopaline synthase activity and kanamycin resistance encoded by the T-DNA insert. A total of 263 T4 plants were grown and assayed

15 for the presence of nopaline in leaf extracts (Errampalli et al., The Plant Cell (1991) 3:149-157).

In addition, T5 seeds were collected from each of the T4 plants and samples of 10-50 seeds were taken to determine the seed fatty acid composition and to

20 determine their ability to germinate in the presence of kanamycin (Feldmann, et al., (1989) Science 243:1351-1354). The 263 plants fell into 3 classes as in Table 5.

2025 SEP 26 1991

TABLE 5

<u>Number of Individuals</u>	<u>Phenotype</u>
63	T4 plants: little or no nopaline present; T5 seeds: wild type fatty acid composition, all kanamycin sensitive
134	T4 plants: nopaline present; T5 seeds: heterozygous fatty acid composition similar to 3707 T3 pool, segregating for kanamycin resistance
64	T4 plants: nopaline present; T5 seeds homozygous mutant fatty acid composition, all kanamycin resistant

The cosegregation of the fatty acid phenotype with the phenotypes conferred by T-DNA sequences in an approximately 1:2:1 pattern provided strong evidence

- 5 that the mutation in line 3707 was the result of a T-DNA insertion. Further experiments were then conducted with the intent of using probes containing T-DNA sequences to clone the T-DNA insert and flanking genomic DNA from line 3707.

10 Preparation of Genomic DNA from Homozygous 3707 Plants

- Seeds from a homozygous line derived from Arabidopsis thaliana (geographic race Wassilewskija (WS)) line 3707 were surface sterilized for 5 min at room temperature in a solution of 5.25% sodium
- 15 hypochlorite (w/v)/0.15% Tween 20 (v/v), then washed several times in sterile distilled water, with a final rinse in 50% ethanol. Immediately following the ethanol wash, the seeds were transferred to sterile filter paper to dry. One to three seeds were then transferred to
- 20 250-mL flasks containing 50 mL of sterile Gamborgs B5 media (Gibco, 500-1153EA), pH 6.0. Cultures were incubated at 22°C, 70 $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{sec}^{-1}$ of continuous light for approximately three weeks, after which time the root tissue was harvested, made into 10 g aliquots (wet
- 25 weight), lyophilized, and stored at -20°C.

36622T 5266T 26B

Using a variation of the procedure of Shure et al., (Cell (1983) 35:225-233) genomic DNA was isolated from the root tissue. Two aliquots of lyophilized tissue were ground to a fine powder using a mortar and pestle.

5 The ground tissue was added to a flask containing 85 mL of lysis buffer (7 M urea, 0.35 M NaCl, 0.05 M Tris-HCl, pH 8.0, 0.02 M EDTA, 1% Sarkosyl, 5% phenol) and mixed gently with a glass rod to obtain a homogeneous suspension. To this suspension an equal volume of

10 phenol:chloroform:isoamyl alcohol (25:24:1) (equilibrated with 10 mM Tris, pH 8, 1 mM EDTA) was added. After the addition of 8.5 mL of 10% SDS the mixture was swirled on a rotating platform for 15 min at room temperature. After centrifugation at 2000xg for 15

15 min, the upper aqueous phase was removed to a new tube and extracted two more times, as above, but without the addition of SDS. To the final aqueous phase was added 1/20th the volume of 3 M potassium acetate, pH 5.5 and two times the volume of ice cold 100% ethanol.

20 Precipitation of the DNA was facilitated by incubation at -20°C for one hour followed by centrifugation at 12,000xg for 10 min. The resulting pellet was resuspended in 3 mL of 10 mM Tris, pH 8, 1 mM EDTA to which was added 0.95 g of cesium chloride (CsCl) and

25 21.4 µL of 10 mg/mL ethidium bromide (EtBr) per mL of solution. The DNA was then purified by centrifugation to equilibrium in a CsCl/EtBr density gradient for 16 h at 15°C, 265,000xg. After removal from the gradient, the DNA was extracted with isopropanol saturated with TE

30 buffer (10 mM Tris, pH 8; 1 mM EDTA) and CsCl to remove EtBr and then dialyzed overnight at 4°C against 10 mM Tris, pH 8, 1 mM EDTA to remove CsCl. The DNA was removed from dialysis and the concentration was determined using the Hoechst fluorometric assay in which

35 an aliquot of DNA is added to 3 mL of 1.5×10^{-6} M bis-

benzimidazole (Hoechst 33258, Sigma) in 1X SSC (0.15 M NaCl, 0.015 M sodium citrate), pH 7.0, incubated at room temperature for 5 min, and read on a fluorometer at excitation 360, emission 450, against a known set of DNA standards.

Plasmid Rescue and Analysis

Five micrograms of genomic DNA from the homozygous 3707 mutant, prepared as described above, was digested with 20 units of either Bam HI or Sal I restriction enzyme (Bethesda Research Laboratory) in a 50 μ L reaction volume according to the manufacturer's specifications. After digestion the DNA was extracted with buffer-saturated phenol (Bethesda Research Laboratory) followed by precipitation in ethanol. The resulting pellet was resuspended in a final volume of 10 μ L of 10 mM Tris, pH 8, and the concentration of the DNA was determined using the Hoechst fluorometric assay as above.

To facilitate circularization, as opposed to end-to-end joining, a dilute ligation reaction was set up containing 250 ng of Bam HI or Sal I digested genomic DNA, 3 Weiss units of T4 DNA ligase (Promega), 50 μ L of 10X ligase buffer (30 mM Tris-HCl, pH 7.8, 100 mM MgCl₂, 100 mM DTT, 5 mM ATP) and 5 μ L of 100 mM ATP in a 500 μ L reaction volume. The reaction was incubated for 16 h at 16°C, heated for 10 min at 70°C, and extracted once with buffer saturated phenol (Bethesda Research Laboratory). The DNA was then precipitated with the addition of two volumes of 100% ethanol and 1/10th volume of 7.5 M ammonium acetate. The resulting pellet was resuspended in a final volume of 10 μ L of 10 mM Tris, pH 8, and the concentration of the DNA was determined using the Hoechst fluorometric assay as above.

Competent DH10B cells (Bethesda Research Laboratory) were transfected with 50 ng of ligated DNA

09210935-122360

at a concentration of 10 ng of DNA per 100 μ L of cells according to the manufacturer's specifications. Transformants from Sal I or Bam HI digests were selected on LB plates (10 g Bacto-tryptone, 5 g Bacto-yeast extract, 5 g NaCl, 15 g agar per liter, pH 7.4) containing 100 μ g/mL ampicillin or 25 μ g/mL kanamycin sulfate, respectively. Ampicillin-resistant (Amp^r ; ampicillin sensitivity, Amp^s) Sal I transformants were screened for the presence of the kanamycin resistance (Kan^r ; kanamycin sensitivity, Kan^s) gene by picking primary transformants and stabbing them first to LB plates containing 100 μ g/mL ampicillin then to LB plates containing 25 μ g/mL kanamycin. After overnight incubation at 37°C the plates were scored for Amp^r/Kan^s colonies. Kanamycin-resistant Bam HI transformants were screened for the presence of the ampicillin resistance gene by picking primary transformants and stabbing them first to LB plates containing 25 μ g/mL kanamycin and then to LB plates containing 100 μ g/mL ampicillin. After overnight incubation at 37°C the plates were scored for Kan^r/Amp^r colonies.

Cultures were made of 192 Amp^r/Kan^s Sal I transformants and 85 Kan^r/Amp^r Bam HI transformants directly into deep-well microtiter plates containing 200 μ L of LB broth (10 g Bacto-tryptone, 5 g Bacto-yeast extract, 5 g NaCl per liter) with 100 μ g/mL ampicillin. Using the Schleicher and Schuell Minifold I apparatus and Nytran membranes, dot blots were set up, in duplicate, using the following conditions: 50 μ L of culture was diluted into 150 μ L of 5X SSC, the culture was lysed and the DNA denatured by the addition of 150 μ L of 0.5 M NaOH, 1.5 M NaCl solution for 3 min at room temperature, the filter was removed from the apparatus and neutralized in 0.5 M Tris, pH 8, 1.5 M NaCl, the DNA was then UV cross-linked to the filters

using the Stratagene Stratalinker, and the filters were heated for 2 h at 80°C and stored at room temperature.

To determine whether T-DNA was contained within any of the rescued plasmids, the dot blots were probed with portions of the right and left borders of T-DNA. The right border probe consisted of a 2.2 kb Hind III-Dra I fragment of DNA obtained from plasmid H23pKC7 (composed of the 3.2 kb Hind III 23 fragment from Ti plasmid pTiC58 (Lemmers et al., J. Mol. Biol. (1989) 144:353-376) cloned into plasmid vector pKC7 (Maniatis et al., Molecular Cloning, A Laboratory Manual (1982) Cold Spring Harbor Laboratory Press)), and the left border probe consisted of a 2.9 kb Hind III-Eco RI fragment obtained from plasmid H10pKC7 (composed of the 6.5 kb Hind III 10 fragment from Ti plasmid pTiC58 (Lemmers et al., J. Mol. Biol. (1989) 144:353-376) cloned into plasmid vector pKC7 (Maniatis et al., Molecular Cloning, A Laboratory Manual (1982) Cold Spring Harbor Laboratory Press)) using standard digestion, electrophoresis, and electroelution conditions as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed (1989) Cold Spring Harbor Laboratory Press). Final DNA purification was obtained by passage of the eluted DNA over an Elutip-D column (Schleicher and Schuell) using the manufacturer's specifications. Concentration of the DNA was determined using the Hoechst fluorometric assay as above. Approximately 100 ng of each probe was labeled with a[³²P]dCTP using a Random Priming Kit from Bethesda Research Laboratories under conditions recommended by the manufacturer. Labeled probe was separated from unincorporated a[³²P]dCTP by passing the reaction through a Sephadex G-25 spun column under standard conditions as described in Sambrook et al., (Molecular

The filters were pre-hybridized in 150 mL of buffer consisting of 6X SSC, 10X Denhardt's solution, 1% SDS, and 100 µg/mL denatured calf thymus DNA for 16 h at 42°C. The denatured, purified, labeled probe was added to the pre-hybridized filters following transfer of the filters to 50 mL of hybridization buffer consisting of 6X SSC, 1% SDS, 10% dextran sulfate, and 50 µg/mL denatured calf thymus DNA. Following incubation of the filters in the presence of the probe for 16 h at 65°C, the filters were washed twice in 150 mL of 6X SSC, 0.5% SDS, twice in 1X SSC, 1% SDS and once in 0.1X SSC, 1% SDS, all at 65°C. The washed filters were subjected to autoradiography on Kodak XAR-2 film at 80°C overnight.

DNA from the Bam HI candidates was made by the alkaline lysis miniprep procedure of Birnboim et al., (Nuc. Acid Res. (1979) 7:1513-1523), as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press). The plasmid DNA was digested with Eco RI restriction enzyme (Bethesda Research Laboratories) in accordance with the manufacturer's specifications and electrophoresed through a 0.8% agarose gel in 1X TBE buffer (0.089 M Tris-borate, 0.089 M boric acid, 0.002 M EDTA). All of the Bam HI candidates which hybridized with the left border probe of T-DNA had the same Eco RI

restriction pattern, which indicated the presence of 14.2 kb of T-DNA and 1.4 kb of putative plant genomic DNA in these clones.

DNA from Sal I candidates was isolated,
 5 restriction-analyzed using Eco RI, Bam HI and Sal I
 enzymes, and electrophoresed through a 0.8% agarose gel,
 as above. All of the Sal I candidates which hybridized
 with the left border probe of T-DNA included 2.9 kb of
 putative plant DNA. Contained within this 2.9 kb
 10 fragment was a 1.4 kb Bam HI-Eco RI fragment as seen
 with the Bam HI rescued plasmids, suggesting that the
 1.4 kb fragment was a subset of the 2.9 kb fragment and
 that it was adjacent to the left border of the T-DNA at
 its site of insertion into the plant genome. Sequence
 15 analysis of one Sal I candidate (pS1) using a primer
 homologous to the left border sequence of T-DNA,
 revealed that the sequence of pS1 was colinear with the
 sequence of the T-DNA left border (Yadav et al., Proc.
 Natl. Acad. Sci. USA (1982) 79:6322-6326) up to
 20 nucleotide 65, followed by non-T-DNA (putative plant)
 sequences.

Southern Analysis with Putative Plant

DNA from Rescued Plasmids

DNA from the seven Bam HI candidates which
 25 hybridized with the left border of the T-DNA was pooled
 and a portion was digested with Eco RI and Bam HI
 restriction endonucleases and electrophoretically
 separated on a 0.8% agarose gel in 1X TBE buffer. After
 excising a 1.4 kb Eco RI-Bam HI fragment from the
 30 agarose gel, the 1.4 kb fragment was purified by use of
 a Gene Clean Kit from Bio 101. Fifty nanograms of the
 resulting DNA fragment was labeled with a[³²P]dCTP using
 a Random Priming Kit (Bethesda Research Laboratory)
 under conditions recommended by the manufacturer.

8622T 966T260

Three micrograms of total genomic DNA from homozygous wild-type Arabidopsis and homozygous 3707 mutant Arabidopsis plants was digested to completion with one of the following restriction enzymes: Sal I, Hind III, Eco RI, Cla I, and Bam HI under conditions suggested by the manufacturer. The digested DNA was subjected to electrophoresis and Southern transfer to Hybond-N membranes (Amersham) as described in Sambrook et al. (Molecular Cloning, A Laboratory Approach, 2nd. ed. (1989) Cold Spring Harbor Laboratory Press). After Southern transfer, the membranes were exposed to UV light using the Stratalinker (Stratagene) as per the manufacturer's instructions, air dried, and heated at 68°C for 2 h.

The filters were prehybridized in 1 M NaCl, 50 mM Tris-Cl, pH 7.5, 1% sodium dodecyl sulfate, 5% dextran sulfate, 100 µg/mL of denatured salmon sperm DNA at 65°C overnight. Fifty nanograms of the radiolabeled 1.4 kb Eco RI-Bam HI plant DNA fragment prepared above was added to the prehybridization solution containing the Southern blot and further incubated at 65°C overnight. The filter was washed for 10 min twice in 200 mL 2X SSPE, 0.1% sodium dodecyl sulfate at 65°C and for 10 min in 200 mL 0.5% SSPE, 0.1% sodium dodecyl sulfate at 65°C. Hybridizing fragments were detected by autoradiography. The analysis confirmed that the probe fragment contained plant DNA and that the T-DNA integration site was in a 2.8 kb Bam HI, a 5.2 kb Hind III, a 3.5 kb Sal I, a 5.5 kb Eco RI, and an approximately 9 kb Cla I fragment of wild type Arabidopsis DNA.

Isolation of Lambda Clones Containing the Wild Type Arabidopsis Delta-15 Desaturase Gene

The 1.4 kb Eco RI-Bam HI fragment (see above) was used as a probe to screen a lGem-11 library made from

genomic DNA isolated from wildtype Arabidopsis thaliana plants, geographic race WS. To construct the library, genomic DNA was partially digested with Sau3A enzyme, and size-fractionated over a salt gradient as described in Sambrook et al. (Molecular Cloning, A Laboratory Approach, 2nd ed. (1989) Cold Spring Harbor Laboratory Press). The size-fractionated DNA was then cloned into Bam HI-digested lGem-11 phage DNA (Promega) following the protocol outlined by the manufacturer. About 25,000 plaque-forming units of phage each were plated on five 150 mm petri plates containing a lawn of KW251 cells on NZY agar media (5 g NaCl, 2 g MgSO₄·7H₂O, 5 g yeast extract, 10 g NZ Amine (casein hydrolysate from ICN Pharmaceuticals), 15 g agar per liter; pH 7.5). The plaques were adsorbed onto nylon membranes (Colony/Plaque Screen, New England Nuclear), in duplicate, and prepared according to the manufacturer's instructions with the addition of a 2 h incubation at 80°C after air drying the filters. The filters were prehybridized at 65°C in hybridization buffer (1% BSA, 0.5 M NaPi, pH 7.2, (NaH₂PO₄ and Na₂HPO₄), 10 mM EDTA, and 7% SDS) for 4 h, after which time they were transferred to fresh buffer containing the denatured radiolabeled probe (see above) and incubated overnight at 65°C. The filters were rinsed twice with 0.1X SSC, 1% SDS at 65°C for 30 min each and subjected to autoradiography on Kodak XA-R film at 80°C overnight. Seven positively-hybridizing plaques were subjected to plaque purification as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press).

Small scale (5 mL) liquid lysates from each of the 7 clones were prepared and titered on KW251 bacteria as described in Sambrook et al. (Molecular Cloning, A Laboratory Manual, 2nd ed (1989), Cold Spring Harbor

Laboratory Press). Phage DNA was isolated using a variation of the method of Chisholm (Biotechniques (1989) 7:21-23) in which the initial lysate was made according to Sambrook et al. (Molecular Cloning, A Laboratory Manual, 2nd ed (1989), Cold Spring Harbor Laboratory Press) the concentration of DNase I and RNase I (Sigma) was reduced by half, and the PEG precipitation step was increased to 16 h. Based on restriction analysis using Hind III, Sal I and Xho I enzymes, the original 7 positive phage fell into 5 different classes. While the average insert size was approximately 15 kb, taken together the clones spanned a 40 kb region of genomic DNA. Through restriction mapping using 4 different enzymes (Hind III, Bam HI, Kpn I, and Sal I) singly, and in pair-wise combinations, accompanied by Southern analysis with the 1.4 kb Eco RI-Bam HI probe (as above) and other probes obtained from the 1 clones themselves, a partial map was obtained in which all 5 clones (11111, 141A1, 14211, 14311 and 14411) were found to share an approximately 3 kb region of homology near the site of T-DNA insertion. Via restriction and Southern analysis, Applicants ascertained that a 5.2 kb Hind III fragment present in clones 1111, 41A1, and 4411 also spanned the site of the T-DNA insertion. This fragment was excised from lambda clone 41A1, inserted into the Hind III site of the pBluescript vector (Stratagene), and the resulting plasmid, designated pF1, was prepared and isolated using standard protocols. This Hind III fragment was subsequently used to probe an Arabidopsis cDNA library (see below).

EXAMPLE 2

CLONING OF ARABIDOPSIS THALIANA DELTA-15
DESATURASE cDNA USING GENOMIC DNA FLANKING
THE T-DNA SITE OF INSERTION IN ARABIDOPSIS THALIANA
5 MUTANT LINE 3707 AS A HYBRIDIZATION PROBE

The 5.2 kb Hind III fragment from plasmid pF1 was purified by electrophoresis in agarose after digestion of the plasmid with Hind III and radiolabeled with ³²p as described above. For the preparation of an
10 Arabidopsis cDNA library, polyadenylated mRNA was prepared from 3 day-old, etiolated Arabidopsis (ecotype Columbia) seedling hypocotyls using standard protocols (Sambrook, et al., Molecular Cloning: A Laboratory Manual, 2nd Ed. (1989) Cold Spring Harbor Laboratory
15 Press). Five micrograms of this mRNA were used as template with an oligo d(T) primer, and Moloney Murine Leukemia Virus reverse transcriptase (Pharmacia) was used to catalyze first strand cDNA synthesis. Second-strand cDNA was made according to Gubler et al., (Gene
20 (1983) 25:263-272) except that DNA ligase was omitted. After the second strand synthesis, the ends of the cDNA were made blunt by reaction with the Klenow fragment of DNA polymerase and ligated to Eco RI/Not I adaptors (Pharmacia). The cDNA's were purified by spun-column
25 chromatography using Sephacryl S-300 and size-fractionated on a 1% low melting point agarose gel. Size-selected cDNAs (1-3 kb) were removed from the gel using agarase (New England Biolabs) and purified by phenol:chloroform extraction and ethanol precipitation.
30 One hundred nanograms of the cDNA was co-precipitated with 1 µg of 1 ZAP II (Stratagene) Eco RI-digested, dephosphorylated arms. The DNAs were ligated in a volume of 4 µL overnight, and the ligation mix was packaged in vitro using the Gigapack II Gold packaging
35 extract (Stratagene).

Approximately 80,000 phage were screened for positively hybridizing plaques using the radiolabeled 5.2 kb Hind III fragment as a probe essentially as described above and in Sambrook et al., (Molecular Cloning: A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press). Replica filters of the phage plaques were soaked in 1 M NaCl, 50 mM Tris-HCl, pH 7.5, 1% SDS, 5% dextran sulfate, 0.1 mg/mL denatured salmon sperm DNA during the pre-hybridization step (8 hr at 65°C) and then probe was added and the hybridization proceeded over 16 hr at the same temperature. Filters were washed sequentially with 2X SSPE, 0.1% SDS at room temperature for 5 min and then again with fresh solution for 10 min, and finally with 0.5X SSPE, 0.1% SDS at 65°C for 5 min. Approximately 20 positively hybridizing plaques were identified in the primary screen. Four of these were picked and subjected to two further rounds of screening and purification. From the tertiary screen, four pure phage plaques were isolated. Plasmid clones containing the cDNA inserts were obtained through the use of a helper phage according to the in vivo excision protocol provided by Stratagene. Double-stranded DNA was prepared using the alkaline lysis method as previously described, and the resulting plasmids were size-analyzed by electrophoresis in agarose gels. The largest one of these, designated pCF3, contained an approximately 1.4 kb insert which was sequenced using Sequenase T7 DNA polymerase (US Biochemical Corp.) and the manufacturer's instructions, beginning with primers homologous to vector sequences that flank the cDNA insert and continuing serially with primers designed from the newly acquired sequences as the sequencing experiment progressed. The sequence of this insert is shown in SEQ ID NO:1.

EXAMPLE 3CLONING OF AN ARABIDOPSIS CDNA ENCODING A PLASTID
DELTA-15 FATTY ACID DESATURASE

5 A related fatty acid desaturase was cloned in a similar fashion, except that the probe used was not derived from a PCR reaction on pCF3, but rather was the actual 1.4 kb Not I fragment isolated from pCF3 which was purified and radiolabeled as described above.

10 Approximately 80,000 phage from the Arabidopsis etiolated hypocotyl cDNA library described above were plated out and screened essentially as before, except as indicated below. The filters were soaked in 1 M NaCl, 50 mM Tris-HCl, pH 7.5, 1% SDS, 5% dextran sulfate, 0.1 mg/mL denatured salmon sperm DNA during the pre-
15 hybridization step (8 hr at 50°C). Then probe was added and the hybridization proceeded over 16 hr at the same temperature. Filters were washed sequentially with 2X SSPE, 0.1% SDS at room temperature for 5 min and then again with fresh solution for 10 min, and finally with
20 0.5X SSPE, 0.1% SDS at 50°C for 5 min. Approximately 17 strongly hybridizing and 17 weakly hybridizing plaques were identified in the primary screen. Four of the weakly hybridizing plaques were picked and subjected to one to two further rounds of screening with the
25 radiolabeled probe as above until they were pure. To ensure that these were not delta-15 desaturase clones, they were further analyzed to determine whether they hybridized to a delta-15 desaturase 3' end-specific probe. The probe used was an 18 bp oligonucleotide
30 which is complementary in sequence (i.e., antisense) to nucleotides 1229 - 1246 of SEQ ID NO:1. The probe was radiolabeled with gamma-³²P ATP using T4 polynucleotide kinase and hybridized to filters containing DNA from the isolated clones in 6X SSC, 5X Denhardt's, 0.1 mg/mL
35 denatured salmon sperm DNA, 1 mM EDTA, 1% SDS at 44°C

B6622T SE661260

overnight. The filters were washed twice in 6X SSC, 0.1% SDS for 5 min at room temperature, then in 6X SSC, 0.1% SDS at 44°C for 3-5 min. After autoradiography of the filters, one of the clones failed to show

5 hybridization to this probe. This clone was picked, and a plasmid clone containing the cDNA insert was obtained through the use of a helper phage according to the in vivo excision protocol provided by Stratagene. Double-stranded DNA was prepared using the alkaline lysis

10 method as previously described, and the resulting plasmid was size-analyzed by electrophoresis in agarose gels following either Not I digestion or digestion with both Nco I and Bgl II. The results were consistent with the presence in this plasmid, designated pCM2, of an

15 approximately 1.3 kb cDNA insert which lacked a 0.7 kb Nco I - Bgl II fragment characteristic of the Arabidopsis delta-15 desaturase cDNA of pCF3. (This fragment corresponds to the DNA located between the Nco I site at nucleotides 474-479 and the Bgl II site at

20 nucleotides 1164-1169 in SEQ ID NO:1). The complete nucleotide sequence of pCM2 is shown in SEQ ID NO:4.

EXAMPLE 4

CLONING OF PLANT FATTY ACID DESATURASE cDNAs FROM OTHER SPECIES BY HYBRIDIZATION TECHNIQUES

25 An approximately 1.4 kb fragment containing the Arabidopsis delta-15 desaturase coding sequence of SEQ ID NO:1 was obtained from plasmid pCF3 through the use of the polymerase chain reaction (PCR). Primers (M13(-20) and T7-17mer primers, 1991 Stratagene

30 Catalogue numbers 300303 and 300302, respectively) flanking the pCF3 insert were used in the PCR which was carried out essentially as described in the instructions provided by the vendor in the Perkin-Elmer/Cetus PCR kit. This fragment was digested with Not I to remove

35 vector sequences, purified by agarose gel electro-

0019931260

phoresis, and radiolabeled with ^{32}P as previously described.

EXAMPLE 5

CLONING OF BRASSICA NAPUS SEED cDNAs ENCODING DELTA-15 FATTY ACID DESATURASES

5 A cDNA library from developing Brassica napus seeds was constructed using the polyadenylated mRNA fraction contained in a polysomal RNA preparation from developing Brassica napus seeds. Polysomal RNA was isolated
10 following the procedure of Kamalay et al., (Cell (1980) 19:935-946) from seeds 20-21 days after pollination. The polyadenylated mRNA fraction was obtained by affinity chromatography on oligo-dT cellulose (Aviv et al., Proc. Natl. Acad. Sci. USA (1972) 69:1408-1411).
15 Four micrograms of polyadenylated mRNA were reverse transcribed and used to construct a cDNA library in lambda phage (Uni-ZAP™ XR vector) using the protocol described in the ZAP-cDNA™ Synthesis Kit (1991 Stratagene Catalog, Item # 200400).
20 For the purpose of cloning the Brassica napus seed cDNAs encoding delta-15 fatty acid desaturases, the Brassica napus seed cDNA library was screened several times using the inserts from the Arabidopsis cDNAs pCF3 and pCM2 as radiolabelled hybridization probes. One of
25 the Brassica napus cDNAs obtained in these screens was used as hybridization probe in a subsequent screen.

For each screening experiment approximately 300,000 phages were screened under low stringency hybridization conditions. The filter hybridizations were carried out
30 in 50 mM Tris pH 7.6, 6X SSC, 5X Denhardt's, 0.5% SDS, 100 ug denatured calf thymus DNA at 50°C overnight and the post hybridization washes were performed in 6X SSC, 0.5% SDS at room temperature for 15 min, then repeated with 2X SSC, 0.5% SDS at 45°C for 30 min, and then

Using the Arabidopsis cDNA insert of pCM2 as a probe in a low stringency screen five strongly hybridizing phages were identified. These phages were purified and excised according to the protocols described in the ZAP-cDNA™ Synthesis Kit and pBluescript II Phagemid Kit (1991 Stratagene Catalog, Item # 200400 and 212205). One of these, designated pBNSF3-f2, contained a 1.3 kb insert. pBNSF3-f2 insert was sequenced completely on both strands. pBNSF3-f2 nucleotide sequence is shown in SEQ ID NO:6. A comparison of this sequence with that of the Arabidopsis thaliana delta-15 desaturase clone (SEQ ID NO:1) confirmed that pBNSF3-f2 is a Brassica napus cDNA that encodes a seed microsomal delta-15 desaturase.

An additional low stringency screen of the Brassica napus seed cDNA library using the cDNA insert in pCM2 as a probe identified eight strongly-hybridizing phages. These phages were plaque purified and used to excise the phagemids as described above. One of these, designated pBNSFd-8, contained a 0.3kb insert. pBNSFd-8 was sequenced completely on one strand, this sequence had significant divergence from the sequence of pBNSF3-f2. The cDNA insert in pBNSFd-8 was used as a hybridization probe in a high stringency screen of the Brassica napus seed cDNA library. The filter hybridizations were carried out in 50 mM Tris pH 7.6, 6X SSC, 5X Denhardt's, 0.5% SDS, 100 ug denatured calf thymus DNA overnight at 50°C and post hybridization washes were in 6X SSC, 0.5% SDS at room temperature for 15 min, then with 2X SSC, 0.5% SDS at 45°C for 30 min, and then twice with 0.2X SSC, 0.5% SDS at 60°C for 30 min. The high stringency screen resulted in three strongly hybridizing phages that were purified and excised as above. One of the

excised plasmids pBNSFd-3 contained a 1.4kb insert that was sequenced completely on both strands. SEQ ID NO:8 shows the nucleotide sequence of pBNSFd-3. A comparison of this sequence with that of the Arabidopsis thaliana delta-15 desaturase clone (SEQ ID NO:4) confirmed that pBNSFd-3 is a Brassica napus cDNA that encodes a seed plastid delta-15 desaturase.

Cloning of a Soybean Seed cDNA Encoding a
Microsomal Delta-15 Glycerolipid Desaturase

10 A cDNA library was made as follows: Soybean embryos (ca. 50 mg fresh weight each) were removed from the pods and frozen in liquid nitrogen. The frozen embryos were ground to a fine powder in the presence of liquid nitrogen and then extracted by Polytron

15 homogenization and fractionated to enrich for total RNA by the method of Chirgwin et al. (Biochemistry (1979) 18:5294-5299). The nucleic acid fraction was enriched for poly A⁺RNA by passing total RNA through an oligo-dT cellulose column and eluting the poly A⁺RNA with salt as

20 described by Goodman et al. (Meth. Enzymol. (1979) 68:75-90). cDNA was synthesized from the purified poly A⁺RNA using cDNA Synthesis System (Bethesda Research Laboratory) and the manufacturer's instructions. The resultant double-stranded DNA was methylated by Eco RI

25 DNA methylase (Promega) prior to filling-in its ends with T4 DNA polymerase (Bethesda Research Laboratory) and blunt-end ligation to phosphorylated Eco RI linkers using T4 DNA ligase (Pharmacia). The double-stranded DNA was digested with Eco RI enzyme, separated from

30 excess linkers by passage through a gel filtration column (Sephacrose CL-4B); and ligated to lambda ZAP vector (Stratagene) according to manufacturer's instructions. Ligated DNA was packaged into phage using the Gigapack packaging extract (Stratagene) according to

35 manufacturer's instructions. The resultant cDNA library

001935 1223

was amplified as per Stratagene's instructions and stored at -80°C.

Following the instructions in the Lambda ZAP Cloning Kit Manual (Stratagene), the cDNA phage library was used to infect *E. coli* BB4 cells and approximately 80,000 plaque forming units were plated onto 150 mm diameter petri plates. Duplicate lifts of the plates were made onto nitrocellulose filters (Schleicher & Schuell). The filters were prehybridized in 25 mL of hybridization buffer consisting of 50mM Tris-HCl, pH 7.5, 1 M NaCl, 1% SDS, 5% dextran sulfate and 0.1 mg/mL denatured salmon sperm DNA (Sigma Chemical Co.) at 50°C for 2 h. Radiolabeled probe prepared from pCF3 as described above was added, and allowed to hybridize for 18 h at 50°C. The probes were washed twice at room temperature with 2X SSPE, 1% SDS for five minutes followed by washing for 5 min at 50°C in 0.2X SSPE, 1% SDS. Autoradiography of the filters indicated that there was one strongly hybridizing plaque, and approximately five weakly hybridizing plaques. The more strongly hybridizing plaque was subjected to a second round of screening as before, excepting that the final wash was for 5 min at 60°C in 0.2X SSPE, 1% SDS. Numerous, strongly hybridizing plaques were observed, and one, well-isolated from other phage, was picked for further analysis.

Following the Lambda ZAP Cloning Kit Instruction Manual (Stratagene), sequences of the pBluescript vector, including the cDNA inserts, from the purified phage was excised in the presence of a helper phage and the resultant phagemid was used to infect *E. coli* XL-1 Blue cells. DNA from the plasmid, designated pXF1, was made by the alkaline lysis miniprep procedure described in Sambrook et al. (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory

Press). The alkali-denatured double-stranded DNA from pXF1 was completely sequenced on both strands. The insert of pXF1 contained a stretch of 1783 nucleotides which contained an unknown open-reading frame and also contained a poly-A stretch of 16 nucleotides 3' to the open reading frame, from nucleotides 1767 to 1783, followed by an Eco RI restriction site. The 2184 bases that followed this Eco RI site contained a 1145 bp open reading frame which encoded a polypeptide of about 68% identity to, and colinear with, the Arabidopsis delta-15 desaturase polypeptide listed in SEQ ID No:2. The putative start methionine of the 1145 bp open-reading frame corresponded to the start methionine of the Arabidopsis microsomal delta-15 peptide and there were no amino acids corresponding to a plastid transit peptide 5' to this methionine. When the insert in pXF1 was digested with Eco RI four fragments were observed, fragments of approximately 370 bp and 1400 bp fragments, derived from the first 1783 bp of the insert in pXF1, and fragments of approximately 600 bp and 1600 bp derived from the the other 2184 nucleotides of the insert in pXF1. Only the 600 bp and 1600 bp fragments hybridized with probe derived from pCF3 on Southern blots. It was deduced that pXF1 contained two different cDNA inserts separated by an Eco RI site and the second of these inserts was a 2184 bp cDNA encoding a soybean microsomal delta-15 desaturase. The complete nucleotide sequence of the 2184 bp soybean microsomal delta-15 cDNA contained in plasmid pXF1 is listed in SEQ ID No:10.

30 Cloning of a Soybean Seed cDNA Encoding a Plastid
 Delta-15 Glycerolipid Desaturase Using
 Soybean Microsomal Delta-15 Desaturase cDNA
 as an Hybridization Probe

35 A 1.0 kb fragment of the coding region of the
 soybean microsomal delta-15 desaturase cDNA contained in

362221 5667260

plasmid pXF1 was excised by digestion with the restriction enzyme Hha I. This 1.0 Kb fragment was purified by agarose gel electrophoresis and radiolabeled with 32P as previously described. The radiolabeled fragment was used to screen 100,000 plaque-forming units of the soybean cDNA library as described above. Autoradiography of the filters indicated that there were eight hybridizing plaques and these were subjected to a second round of screening. Sequences of the pBluescript vector from all eight of the purified phages, including the cDNA inserts, were excised in the presence of a helper phage and the resultant phagemids were used to infect *E. coli* XL-1 Blue cells. DNA from the plasmids was made by the alkaline lysis miniprep procedure described in Sambrook et al. (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press). Restriction analysis showed they contained inserts ranging from 1.0 kb to 3.0 kb in size. One of these inserts, designated pSFD-118bwp, contained an insert of about 1700 bp. The alkali-denatured double-stranded DNA from pSFD-118bwp was completely sequenced on both strands. The insert of pSFD-118bwp contained a stretch of 1675 nucleotides which contained an open-reading frame encoding a polypeptide of about 80% identity with, and colinear with, the Arabidopsis plastid delta-15 desaturase polypeptide listed in SEQ ID No:5. The open-reading frame also encoded amino acids corresponding to a plastid transit peptide at the 5' end of the open-reading frame. The transit peptide was colinear with, and shared some homology to, the transit peptide described for the *Arabidopsis* plastid delta-15 glycerolipid desaturase. Based on the homology to *Arabidopsis* plastid delta-15 glycerolipid desaturase and because of the presence of a plastid transit peptide, the cDNA contained in plasmid pSFD-118bwp was deduced to

be a soybean plastid delta-15 glycerolipid desaturase. The complete nucleotide sequence of the 1675 bp soybean plastid delta-15 glycerolipid desaturase cDNA is listed in SEQ ID NO:12.

5

EXAMPLE 6

CLONING OF cDNA SEQUENCES ENCODING FATTY ACID
DESATURASES BY POLYMERASE CHAIN REACTION

Analysis of the deduced protein sequences of the different higher plant glycerolipid desaturases described in this invention reveals to those skilled in the art regions of the amino acid sequences that have been conserved among higher plants and between higher plants and cyanobacterial des A. These short stretches of amino acids can be used to design oligomers as primers for polymerase chain reactions. Two amino acid sequences that are highly conserved between the des A and plant delta-15 desaturases polypeptides are amino acid sequences 97-108 and 299-311 (SEQ ID NO:2). Polymerase chain reactions (PCRs) were performed using GeneAmp® RNA PCR Kit (Perkin Elmer Cetus) following manufacturer's protocols. In one PCR experiment, SEQ ID NOS:22 and 23 were used as sense primers and either SEQ ID NOS:24 and 25 or SEQ ID NOS:26 and 27 as antisense primers on poly A+ RNA purified from both Arabidopsis leaf and canola developing seeds. For this, ca. 100 ng of polyA+ RNA was isolated as described previously and reverse-transcribed using the kit using random hexamers. Then the cDNA was used in PCR using 64 pmoles each of SEQ ID NOS:22 and 23 as sense primers and either a mixture of 64 pmoles of SEQ ID NO:24 and 78 pmoles of SEQ ID NO:25 or a mixture 35 pmoles of SEQ ID NO:26 and 50 pmoles of SEQ ID NO:27 by the following program: a) 1 cycle of 2 min at 95°C and 15 C at 50°C, b) 30 cycles of 3 min at 65°C (extension), 1 min 20 sec at 95°C (denaturation), 2 min at 50°C (annealing), and c) 1

8621993 122399

cycle of 7 min at 65°C. PCR products were analyzed by gel electrophoresis. All PCRs resulted in PCR products of the correct size (ca. 630 bp). The PCR products from Arabidopsis and canola were purified and used as

5 radiolabeled hybridization probes to screen the Lambda Yes Arabidopsis cDNA library at low stringency, as described above. This led to the isolation of a pure phage, which was excised to give plasmid pYacp7. The cDNA insert in pYacp7 was partially sequenced. Its

10 sequence showed that it encoded an incomplete desaturase polypeptide that was identical to another cDNA (in plasmid pFadx-2) isolated by low-stringency hybridization as described previously. The composite sequence derived from the partial sequences from the

15 cDNA inserts in pFadx-2 and pYacp7 is shown in SEQ ID NO:16 and the polypeptide encoded by it in SEQ ID NO:17. As discussed previously, SEQ ID NO:17 is a putative plastid delta-15 desaturase. A full-length version of pYacp7 can be readily isolated using it has a

20 hybridization probe.

Two additional conserved regions correspond to aminoacid residues 130 to 137 and 249 and 256 of SEQ ID NO:7 (*Brassica napus* glycerolipid desaturase delta-15). Degenerate oligomers were designed to these regions with

25 additional nucleotides containing a restriction site for Bam H1 were added to the 5' ends of each oligonucleotide to facilitate subcloning of the PCR products. The nucleotide sequences of these oligonucleotides named F2-3 and F2-3c are shown in SEQ ID NO:18 and SEQ ID

30 NO:19 respectively.

Mixtures of degenerate oligonucleotides F2-3 and F2-3c were used to amplify, isolate and clone glycerolipid desaturase sequences represented in corn seed mRNA population, essentially as described in the GeneAmp RNA

35 PCR Kit purchased from Perkin Elmer Cetus and in Innis,

et al., Eds, (1990) PCR Protocols: A Guide to Methods and Applications, Academic Press, San Diego.

Corn seed RNA was obtained from developing corn seeds 15-20 days after pollination by the method of Chirgwin et al., (1979) Biochemistry 18:5294. Corn seed polyadenylated mRNA was isolated by affinity chromatography on oligo-dT cellulose (Aviv et al., Proc. Natl. Acad. Sci. USA (1972) 69:1408-1411). 20-50ng of A+mRNA were used in reverse transcription reactions with oligo-dT and random hexamers primers using the reaction buffer and conditions recommended by Perkin Elmer Cetus. The resulting cDNA was then used as template for the amplification of corn seed glycerolipid sequences using the set of degenerate primers in SEQ ID NO: 18 and 19. Reaction conditions were as described by Perkin Elmer Cetus, the amplification protocol consisted of a sequence of 95°C/1 min, 55°C/1 min, 72°C/2 min for 30-50 cycles. The resulting polymerase reaction products were phenol-chloroform extracted, digested with Bam HI and separated from unincorporated primers by gel filtration chromatography on Linker 6 spin columns (Pharmacia Inc.). The resulting PCR products were cloned into pBluescript SK at the Bam HI site, and transformed into *E. coli* DH5 competent cells. Restriction analysis of plasmid DNA from the transformed colonies obtained revealed a colony, PCR-20, that contained an insert of about 0.5 kB in size at the pBluescript SK BamHI site. The PCR-20 insert was completely sequenced on both strands. The nucleotide sequence of PCR20 insert is shown in SEQ ID NO:14 and the translated amino acid sequence is shown in SEQ ID NO:15. This amino acid sequence shows an overall identity of 61.9% to the amino acid sequence of Brassica napus microsomal delta-15 deaturase shown in SEQ ID NO:7. This result identifies the PCR20 insert as a polymerase reaction product of a

corn seed delta-15 desaturase cDNA. PCR20 insert may be used as a probe to readily isolate full length corn seed delta-15 desaturase cDNAs or as such to antisense or cosuppress corn seed glycerolipid delta-15 desaturase gene expression in transgenic corn plants by cloning it in the appropriate corn gene expression vector.

EXAMPLE 7

USE OF THE ARABIDOPSIS THALIANA DELTA-15 DESATURASE GENOMIC CLONES AS A RESTRICTION FRAGMENT LENGTH POLYMORPHISM (RFLP) MARKERS TO MAP THE DELTA-15 DESATURASE LOCI IN ARABIDOPSIS.

DNA flanking the T-DNA insertion site in mutant line 3707 was used to map the genetic locus encoding the delta-15 desaturase of Arabidopsis thaliana seeds. An approximately 12 kB genomic DNA fragment containing the Arabidopsis delta-15 desaturase coding sequence was removed from the lambda-4211 clone by digestion with restriction endonuclease Xho I, separated from the Lambda arms by agarose gel electrophoresis, and purified using standard procedures. The isolated DNA was labeled with ³²P using a random priming kit from Pharmacia under conditions recommended by the manufacturer. The radioactive DNA was used to probe a Southern blot containing genomic DNA from Arabidopsis thaliana (ecotype Wassileskija and marker line W100 ecotype Landesberg background) digested with one of several restriction endonucleases. Following hybridization and washes under standard conditions (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press), autoradiograms were obtained. Different patterns of hybridization (polymorphisms) were identified in digests using restriction endonucleases Bgl II, Cla I, Hind III, Nsi I, and Xba I. The same radiolabeled DNA fragment was used to map the polymorphism essentially as described by

001993 12298

Helentjaris et al., (Theor. Appl. Genet. (1986) 72:761-769). The radiolabeled DNA fragment was applied as described above to Southern blots of Xba I digested genomic DNA isolated from 117 recombinant inbred progeny (derived from single-seed descent lines to the F₆ generation) resulting from a cross between Arabidopsis thaliana marker line W100 and ecotype Wassileskija (Burr et al., Genetics (1988) 118:519-526). The bands on the autoradiograms were interpreted as resulting from inheritance of either paternal (ecotype Wassileskija) or maternal (marker line W100) DNA or both (a heterozygote). The resulting segregation data were subjected to genetic analysis using the computer program Mapmaker (Lander et al., Genomics (1987) 1:174-181). In conjunction with previously obtained segregation data for 63 anonymous RFLP markers and 9 morphological markers in Arabidopsis thaliana (Chang et al., Proc. Natl. Acad. Sci. USA (1988) 85:6856-6860; Nam et al., Plant Cell (1989) 1:699-705), a single genetic locus was positioned corresponding to the genomic DNA containing the delta-15 desaturase coding sequence. The location of the delta-15 desaturase gene was thus determined to be on chromosome 2 between the lambda AT283 and cosmid c6842 RFLP markers, near the py and erecta morphological markers.

The cDNA in plasmid pCM2 was also shown to hybridize polymorphically to genomic DNA from Arabidopsis thaliana (ecotype Wassileskija and marker line W100 ecotype Landesberg background) digested with Eco RI. It was used as a RFLP marker to map the genetic locus for the gene encoding this fatty acid desaturase in Arabidopsis as described above. A single genetic locus was positioned corresponding to this desaturase cDNA. Its location was thus determined to be on chromosome 3 between the lambda AT228 and cosmid c3838

RFLP markers, "north" of the glabrous locus (Chang et al., Proc. Natl. Acad. Sci. USA (1988) 85:6856-6860; Nam et al., Plant Cell (1989) 1:699-705).

EXAMPLE 8

5 USE OF SOYBEAN SEED MICROSOMAL DELTA-15 GLYCEROLIPID DESATURASE cDNA SEQUENCE IN PLASMID AS A RESTRICTION 10 FRAGMENT LENGTH POLYMORPHISM (RFLP) MARKER

15 A 600 bp fragment of the cDNA insert from plasmid pXF1, which contains about 300 bp of the coding sequence and 300 bp of the 3' untranslated sequence, was excised by digestion with restriction enzyme Eco RI in standard conditions as described in Sambrook et al. (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press), purified by agarose gel electrophoresis and labeled with ³²P using a Random Priming Kit from Bethesda Research Laboratories under conditions recommended by the manufacturer. The resulting radioactive probe was used to probe a Southern blot (Sambrook et al., Molecular Cloning, A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press) containing genomic DNA from soybean [Glycine max (cultivar Bonus) and Glycine soja (PI81762)], digested with one of several restriction enzymes. After hybridization and washes under standard conditions (Sambrook et al. Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press), autoradiograms were obtained and different patterns of hybridization (polymorphisms) were identified in digests performed with restriction enzymes Bam HI, Eco RV and Eco RI. The same probe was then used to map the polymorphic pXF1 locus on the soybean genome, essentially as described by Helentjaris et al. (Theor. Appl. Genet. (1986) 72:761-769). Plasmid pXF1/600 bp probe was applied, as described above, to Southern blots of EcoRI, PstI, EcoRV, BamHI, or Hin DIII digested

BBE22T 506T249

genomic DNAs isolated from 68 F2 progeny plants resulting from a *G. max* Bonus x *G. soja* PI81762 cross. The bands on the autoradiograms were interpreted as resulting from the inheritance of either paternal (Bonus) or maternal (PI81762) pattern, or both (a heterozygote). The resulting data were subjected to genetic analysis using the computer program Mapmaker (Lander et al., Genomics (1987) 1:174-181). In conjunction with previously obtained data for 436 anonymous RFLP markers in soybean (Tingey et al., J. Cell. Biochem., Supplement 14E (1990) p. 291, abstract R153], Applicants were able to position a single genetic locus corresponding to the pXF1/600 bp probe on the soybean genetic map. This confirms that the gene for microsomal delta-15 desaturase is located on chromosome 19 in the soybean genome. This information will be useful in soybean breeding targeted towards developing lines with altered polyunsaturate levels.

20

EXAMPLE 9

OVEREXPRESSION OF MICROSOMAL DELTA-15
FATTY ACID DESATURASE IN PLANTS

Detailed procedures for DNA manipulation, such as use of restriction endonucleases and other DNA modifying enzymes, agarose gel electrophoresis, isolation of DNA from agarose gels, transformation of *E. coli* cells with plasmid DNA, and isolation and sequencing of plasmid DNA are described in Sambrook et al. (1989) Molecular cloning, A Laboratory Manual, 2nd ed. Cold Spring Harbor Laboratory Press and Ausubel et al. (1989) Current Protocols in Molecular Biology John Wiley & Sons. All restriction enzymes and modifying enzymes were obtained from Bethesda Research Laboratory, unless otherwise noted.

To test the biological effect of overexpression of the microsomal delta-15 desaturase SEQ ID NO:1, i.e., the cDNA encoding Arabidopsis thaliana microsomal delta-15 desaturase, was placed in the sense orientation

5 behind either the CaMV 35S promotor, to provide constitutive expression, or behind the promotor for the gene encoding soybean α' subunit of the β -conglycinin (7S) seed storage protein, to provide embryo-specific expression. To create the chimeric gene constructs,

10 specific expression cassettes were made to facilitate easy manipulation of the desired clones. The chimeric genes were then transformed into plant cells by Agrobacterium tumefaciens's binary Ti plasmid vector system [Hoekema et al., (1983) Nature 303:179-180; Bevan

15 (1984) Nucl. Acids Res. 12:8711-8720].

Overexpression of Arabidopsis Delta-15 Fatty Acid
Desaturase in Transgenic Carrot Hairy Roots

To confirm the identity of SEQ ID NO:1 (Arabidopsis microsomal delta-15 fatty acid desaturase) and to test

20 the biological effect of its overexpression in a heterologous plant species, the constitutive chimeric gene 35S:SEQ ID NO:1 was introduced into carrot tissue by Agrobacterium. The cassette for constitutive gene expression in plasmid, pAW28, originated from pK35K

25 which, in turn, is derived from pKNK. Plasmid pKNK is a pBR322-based vector containing a chimeric gene for plant kanamycin resistance: nopaline synthase (NOS) promoter/neomycin phosphotransferase (NPT) II coding region/3' NOS chimeric gene. Plasmid pKNK has been

30 deposited on 7 January 1987 with the American Type Culture Collection of Rockville, Maryland, USA under the provisions of the Budapest Treaty and bears the deposit accession number 67284. A map of this plasmid is shown in Lin, et al., Plant Physiol. (1987) 84:856-861. The

35 NOS promoter region is a 296 bp Sau 3A-Pst I fragment

1993-12-23 09:22:33

corresponding to nucleotides -263 to +33, with respect to the transcription start site, of the NOS gene described by Depicker et al. (1982) J. Appl. Genet. 1:561-574. The Pst I site at the 3' end was created at the translation initiation codon of the NOS gene. The NptII coding region is a 998 bp Hind III-Bam HI fragment obtained from transposon Tn5 (Beck et al., Gene (1982) 19:327-336) by the creation of Hind III and Bam HI sites at nucleotides 1540 and 2518, respectively. The 3' NOS is a 702 bp Bam HI-Cla I fragment from nucleotides 848 to 1550 of the 3' end of the NOS gene (Depicker et al., J. Appl. Genet. (1982) 1:561-574) including its polyadenylation region. pKNK was converted to pK35K by replacing its Eco RI-Hind III fragment containing the NOS promoter with a Eco RI-Hind III fragment containing the CaMV 35S promoter. The Eco RI-Hind III 35S promoter fragment is the same as that contained in pUC35K that has been deposited on 7 January 1987 with the American Type Culture Collection under the provisions of the Budapest Treaty and bears the deposit accession number 67285. The 35S promoter fragment was prepared as follows, and as described in Odell et al., Nature (1985) 313:810-813, except that the 3' end of the fragment includes CaMV sequences to +21 with respect to the transcription start site. A 1.15 KB Bgl II segment of the CaMV genome containing the region between -941 and +208 relative to the 35S transcription start site was cloned in the Bam HI site of the plasmid pUC13. This plasmid was linearized at the Sal I site in the polylinker located 3' to the CaMV fragment and the 3' end of the fragment was shortened by digestion with nuclease Bal31. Following the addition of Hind III linkers, the plasmid DNA was recircularized. From nucleotide sequence analysis of the isolated clones, a 3' deletion fragment was selected with the Hind III

linker positioned at +21. The 35S promoter fragment was isolated as an Eco RI-Hind III fragment, the Eco RI site coming from the polylinker of pUC13.

SECRET
5 The NPTII coding region in plasmid pK35K was removed from plasmid pK35K by digestion with Hind III and Bam HI restriction enzymes. Following digestion, the ends of the DNA molecules were filled-in using Klenow enzyme. Not I linkers (New England Biolabs) were then ligated on the ends and the plasmid was
10 recircularized to yield plasmid pK35Nt. A 1.7 kB fragment containing the 35S promoter region - Not I site - 3' untranslated region from nopaline synthase was liberated from pK35Nt using restriction endonucleases Eco RI and Cla I. Following restriction digestion the
15 ends of the DNA molecules were filled-in using Klenow enzyme after which Xho I linkers (New England Biolabs) were added. The 1.7 kB fragment, now containing Xho I sites at either end, was gel isolated and cloned into the plasmid vector pURA3 (Clontech) at its unique Xho I
20 site. The vector pURA3 was chosen due to the absence of a Not I restriction site, the presence of a single Xho I restriction site and because the relatively large size of the vector (pURA3) would make the isolation of the gene expression cassettes relatively easy from the
25 final construct.

The 1.4 kB Not I fragment in plasmid pCF3 containing Arabidopsis microsomal delta-15 desaturase (SEQ ID NO:1) was isolated and ligated to pAW28 (the constitutive expression cassette) previously linearized
30 with Not I restriction enzyme and treated with calf intestinal alkaline phosphatase (Boehringer Mannheim) to result in plasmids pAW29 and pAW30 that had SEQ ID NO:1 cloned in a sense orientation and antisense orientation, respectively, with respect to the promoter. The
35 orientation of the cDNA relative to the promoters was

established by digestion with appropriate restriction endonucleases or by sequencing across the promotor-cDNA junctions.

The chimeric genes 35S promotor/sense SEQ ID NO:1/3'NOS and 35S promotor/antisense SEQ ID NO:1/3'NOS were isolated as a 3 kB Xho I fragment from plasmids pAW29 and pAW30, respectively, and cloned into the binary vector pZS194b at its unique Sal I site to result in plasmids pAW31 and pAW32, respectively. The orientation of the plant selectable marker gene in pAW31 and pAW32 is the same as that of the 35S promoter as ascertained by digestion with appropriate restriction endonucleases. Binary vector pZS194b contains the pBR322 origin of replication, the replication and stability regions of the Pseudomonas aeruginosa plasmid pVS1 [Itoh, et al., (1984) Plasmid 11:206-220] required for replication and maintenance of the plasmid in Agrobacterium, the bacterial NPT II gene (kanamycin resistance) from Tn5 [Berg et al., (1975) Proc. Nat'l. Acad. Sci. U.S.A. 72:3628-3632] as a selectable marker for transformed bacteria, left and right borders of the T-DNA of the Ti plasmid [Bevan et al., (1984) Nucl. Acids Res. 12:8711-8720], and, between the left and right T-DNA borders are the chimeric NOS:NPT II gene for plant kanamycin resistance, described above, as a selectable marker for transformed plant cells and the E. coli lacZ a-complementing segment [Vieria and Messing (1982) Gene 19:259-267] with unique restriction endonuclease sites for Kpn I and Sal I.

The binary vectors pAW31 and pAW32 were transformed by the freeze/thaw method [Holsters et al. (1978) Mol. Gen. Genet. 163:181-187] into Agrobacterium tumefaciens strain R1000, carrying the Ri plasmid pRiA4b from Agrobacterium rhizogenes [Moore et al., (1979) Plasmid

SECRET 561249

2:617-626] to result in transformants R1000/pAW31 and R1000/pAW32, respectively.

Carrot (Daucus carota L.) cells were transformed by co-cultivation of carrot root disks with strain R1000, R1000/pAW31, or R1000/pAW32 by the method of Petit et al., (1986) Mol. Gen. Genet. 202:388-393]. To prepare explants for inoculation, carrots purchased from the local supermarket were first scrubbed gently with water and dish detergent, then rinsed thoroughly with tap and distilled water. They were surface sterilized in a stirred solution of 50% Clorox and distilled water for 30 min and rinsed thoroughly with sterile distilled water. The carrots were peeled using an autoclaved vegetable peeler and then sliced with a scalpel blade into disks of approximately 5-10 mm thickness. The disks were placed in petri dishes, onto a medium consisting of distilled deionized water solidified with 0.7% agar, in an inverted orientation so that the cut surface nearest to the root apex of the carrot was exposed for inoculation.

Cultures of Agrobacterium strains R1000, R1000/pAW31, and R1000/pAW32 were initiated from freshly grown plates in LB broth plus the appropriate antibiotic selective agents (50 mg/L chloramphenicol for the R1000 or 50 mg/L each of chloramphenicol and kanamycin for R1000/pAW31 and R1000/pAW32) and grown at 28°C to an optical density of around 1.0 at 600 nm. Bacterial cells were pelleted by centrifugation, rinsed and resuspended in LB broth without antibiotics. Freshly cut carrot disks were inoculated by applying 100 µL of the bacterial suspension to the cut surface of each disk. As a control, some disks were inoculated with sterile LB broth only, to indicate the extent of root formation in the absence of Agrobacterium.

Inoculated root disks were incubated at 25°C in the dark in petri dishes sealed with Parafilm. After two weeks of co-cultivation of carrot disks with Agrobacterium, the carrot disks were transferred to fresh agar-solidified water medium containing 500 mg/L carbenicillin for the counterselection of Agrobacterium. At this time, hairy root formation was noted on some root disks. Transfer of the explants to fresh counterselection medium was done at four weeks. Excision of individual roots from the explants was begun at six weeks. Ten days later, additional roots were taken from the explants as needed.

Approximately 5-10 mm long hairy roots were excised and individually subcultured on MS minimal organics medium with 30 g/L sucrose (Gibco, Grand Island, N. Y., Cat. No. 510-1118EA) and 500 mg/L carbenicillin. Approximately equal numbers of roots were subcultured in liquid medium and in a medium solidified with 0.6% agarose. Cultures on solid medium were grown in 60 x 100 mm petri dishes, liquid cultures were in 6-well culture dishes. When excising roots, an effort was made to select single roots from distinct callus-like outgrowths on the wounded surface. These sites of excision were marked on the lid of the petri dish to minimize repeat sampling of tissue originating from the same transformation event.

Two to three weeks after excision from the explants, individual hairy root cultures that were not visibly contaminated with Agrobacterium were transferred to fresh MS medium supplemented with 500 mg/L carbenicillin. The root mass of each culture was cut into segments including one or more branch roots, and these segments were transferred as a group to a plate or well of fresh medium. Approximately 20 mg fresh weight of tissue of root cultures which grew to adequate size

8522T-5E6T260

within the next two to three weeks were sampled for fatty acid composition by gas chromatography of the fatty acyl methyl esters essentially as described by Browse et al., (Anal. Biochem. (1986) 152:141-145) except that 2.5% H₂SO₄ in methanol was used as the methylation reagent and samples were heated for 1.5 h at 80°C to effect the methanolysis of the seed triglycerides. The results are shown in Table 6. A second sample of tissue consisting of an actively growing root tip of approximately 1 cm was excised and placed on MS medium supplemented with 500 mg/L carbenicillin and 25-50 mg/L kanamycin to test for kanamycin resistance select for hairy roots co-transformed with the binary vector [Simpson et al. (1986) Plant Mol. Biol. 6:403-415].

TABLE 6
 Percent 18:3 and 18:2/18:3 Ratio in
Roots of Transgenic Carrots

<u>Root Sample</u>	<u>Transformation Vector Used</u>	<u>%18:3</u>	<u>%18:2/18:3</u>
1	R1000/pAW31	62	0.09
2	R1000/pAW31	8	7.30
3	R1000/pAW31	10	5.69
4	R1000/pAW31	62	0.06
5	R1000/pAW31	10	5.07
6	R1000/pAW31	4	14.2
7	R1000/pAW31	61	0.18
8	R1000/pAW31	4	15.1
9	R1000/pAW31	61	0.07
10	R1000/pAW31	63	0.09
11	R1000/pAW31	15	3.04
12	R1000/pAW31	64	0.14
13	R1000/pAW31	5	9.94
14	R1000/pAW31	9	6.72
15	R1000/pAW31	8	7.08
16	R1000/pAW31	8	6.31
17	R1000/pAW31	23	1.86
18	R1000/pAW31	8	7.33
19	R1000/pAW31	10	5.99
20	R1000/pAW31	7	8.83
21	R1000/pAW32	9	6.80

<u>Root Sample</u>	<u>Transformation Vector Used</u>	<u>%18:3</u>	<u>%18:2/18:3</u>
22	R1000/pAW32	4	11.8
23	R1000/pAW32	3	18.8
24	R1000/pAW32	10	6.21
25	R1000/pAW32	7	8.57
26	R1000/pAW32	3	16.4
27	R1000/pAW32	6	8.29
28	R1000/pAW32	5	9.19
29	R1000/pAW32	5	8.47
30	R1000/pAW32	8	7.17
31	R1000/pAW32	4	11.9
32	R1000/pAW32	8	7.20
33	R1000/pAW32	5	10.4
34	R1000/pAW32	8	7.29
35	R1000/pAW32	3	17.2
36	R1000/pAW32	8	7.27
37	R1000/pAW32	9	6.01
38	R1000/pAW32	9	6.62
40	R1000/pAW32	9	6.02
41	R1000	8	7.23
42	R1000	8	7.83
43	R1000	10	6.20
44	R1000	9	5.97
45	R1000	9	6.73
46	R1000	9	6.27
47	R1000	8	7.27
48	R1000	7	8.30
49	R1000	9	7.11

5 The ability of R1000 transformed "hairy" roots to grow in the absence of exogenous phytohormones can be attributed to the Ri plasmid, pRiA4b. When R1000/pAW31 or R1000/pAW32 strains are used to transform, only a fraction (about half) of the "hairy" roots will also be

transformed with the experimental binary vector, pAW31 or pAW32. Thus, as expected, not all hairy roots resulting from transformation with R1000/pAW31 show the high 18:3 phenotype. The absence of any significant fatty acid phenotype in "hairy roots" transformed with R1000/pAW31 is expected, since carrot and Arabidopsis delta-15 desaturase sequences are not expected to be sufficiently related. These results show that overexpression of Arabidopsis microsomal delta-15 desaturase can result in over 10-fold increase in 18:3 at the expense of 18:2 in heterologous plant tissue.

Overexpression of Arabidopsis Delta-15 Fatty Acid Desaturase in Seeds and Complementation of the Mutation in Delta-15 Desaturation in Mutant 3707

To complement the delta-15 desaturation mutation in the T-DNA mutant 3707 and to test the biological effect of overexpression of SEQ ID NO:1 (Arabidopsis microsomal delta-15 fatty acid desaturase) in seed, the embryo-specific promoter:SEQ ID NO:1 chimeric gene was transformed into the mutant plant. This embryo-specific expression cassette in pAW42 was produced, in part, using a modified version of vector pCW109. Vector pCW109 itself was made by inserting into the Hind III site of the cloning vector pUC18 (Bethesda Research Laboratory) a 555 bp 5' non-coding region (containing the promoter region) of the β -conglycinin gene followed by the multiple cloning sequence containing the restriction endonuclease sites for Nco I, Sma I, Kpn I and Xba I, then 1174 bp of the common bean phaseolin 3' untranslated region into the Hind III site [Slightom et al., Proc. Nat'l Acad. Sci. U.S.A. (1983) 80:1897-1901]. The β -conglycinin promoter region used is an allele of the published β -conglycinin gene (Doyle et al., J. Biol. Chem. (1986) 261:9228-9238) due to differences at 27

nucleotide positions. Further sequence description may be found in Slightom (W091/13993).

The modifications to vector pCW109 were as follows: The potential translation start site was destroyed by digestion with Nco I and Xba I restriction enzymes followed by treatment with mung bean nuclease (New England Biolabs) to create linear, blunt ended DNA molecules. After ligation of Not I linkers (New England Biolabs) and digestion with Not I restriction enzyme (New England Biolabs) the plasmid was recircularized. Confirmation of the desired change was obtained by dideoxy sequencing. The resulting plasmid was designated pAW35. The 1.8 kB Hind III fragment from pAW35 containing the modified β -conclycinin promotor/3' phaseolin region was subcloned into the Hind III site in plasmid vector pBluescript SK⁺ (Stratagene) creating plasmid pAW36. Plasmid pAW36 was linerized at its unique Eco RI site and ligated to Eco RI/Xho I adaptors (Stratagene). Following digestion with Xho I, the 1.7 kB Xho I fragment containing the β -conclycinin promotor/Not I site/3'-phaseolin untranslated region was cloned into the Xho I site in pURA3 vector (Clonetech). The resultant plasmid, pAW42, contains the seed specific expression cassette bordered by Xho I sites to facilitate cloning into modified T-DNA binary vectors and a unique Not I site to facilitate cloning of target cDNA sequences. Vector pURA3 was chosen due to the absence of a Not I restriction site, the presence of a single Xho I restriction site, and the relatively large size of the vector (pURA3) would make the isolation of the gene expression cassettes relatively easy from the final construct.

The 1.4 kB Not I fragment in plasmid pCF3 containing Arabidopsis microsomal delta-15 desaturase (SEQ ID NO:1) was isolated and ligated to plasmid pAW42

09230935 123330

(the seed-specific expression cassette) that had previously been linearized with Not I restriction enzyme and treated with calf intestinal alkaline phosphatase (Boehringer Mannheim) to result in plasmids pAW45 that had SEQ ID NO:1 cloned in a sense orientation with respect to the promoter. The orientation of the cDNA relative to the promoters was established by digestion with appropriate restriction endonucleases or by sequencing across the promotor-cDNA junctions.

The chimeric β -conclycinin promotor/sense SEQ ID NO:1/phaseolin 3' was isolated as a 3.2 kB Xho I fragment from plasmid pAW45 and subcloned into the binary vector pAW25 at its unique Sal I site. In the resulting vector, pAW50, the orientation of the plant selectable marker is the same as that of the β -conclycinin promoter as ascertained by digestion with appropriate restriction endonucleases. Plasmid pAW25, is derived from plasmids pZS94K and pML2. Plasmid pZS94K contains the pBR322 origin of replication, the replication and stability regions of the Pseudomonas aeruginosa plasmid pVS1 [Itoh, et al., (1984) Plasmid 11:206-220] required for replication and maintenance of the plasmid in Agrobacterium, the bacterial NPT II gene (kanamycin resistance) from Tn5 [Berg et al., (1975) Proc. Nat'l. Acad. Sci. U.S.A. 72:3628-3632] as a selectable marker for transformed bacteria, a T-DNA left border fragment of the octopine Ti plasmid pTiA6 and right border fragment derived from TiAch5 describe by van den Elzen et al. (Plant Mol. Biol. (1985) 5:149-154). Between these borders are the E. coli lacZ a-complementing segment [Vieria and Messing (1982) Gene 19:259-267] with restriction endonuclease sites Sal I and Asp 718 derived from pUC18. A 4.5 kB Asp 718-Sal I DNA fragment containing the chimeric herbicide sulfonylurea (SU)-resistant acetolactate (ALS) gene was

obtained from plasmid pML2 and cloned into the Asp
718-Sal I sites of plasmid pZS94K. This chimeric ALS
gene contained the CaMV 35S promoter/Cab22L Bgl II-Nco I
fragment that is described by Harpster et al., [Mol.
5 Gen. Genet. (1988) 212:182-190] and the Arabidopsis ALS
coding and 3' non-coding sequences [Mazur et al., (1987)
Plant Physiol. 85:1110-1117] that was mutated so that it
encodes a SU-resistant form of ALS. The mutation,
introduced by site-directed mutagenesis, are those
10 present in the tobacco SU-resistant Hra gene described
by Lee et al., (1988) EMBO J. 5:1241-1248. The
resulting plasmid was designated pAW25.

The binary vector pAW25 containing the chimeric
embryo-specific β -conglycinin promotor:sense SEQ ID NO:1
15 gene was transformed by the freeze/thaw method [Holsters
et al., (1978) Mol. Gen. Genet. 163:181-187] into the
avirulent Agrobacterium strain LBA4404/pAL4404 [Hoekema
et al., (1983) Nature 303:179-180].

Arabidopsis root cultures were transformed by co-
20 cultivation with Agrobacterium using standard aseptic
techniques for the manipulation of sterile media and
axenic plant/bacterial cultures were followed, including
the use of a laminar flow hood for all transfers.
Compositions of the culture media are listed in Table 8.
25 Unless otherwise indicated, 25x100 mm petri plates were
used for plant tissue cultures. Incubation of plant
tissue cultures was at 23°C under constant illumination
with mixed fluorescent and "Gro and Sho" plant lights
(General Electric) unless otherwise noted. To initiate
30 in vitro root cultures of the T-DNA homogyous mutant
line 3707 (Arabidopsis thaliana (L.) Heynh, geographic
race Wassilewshija) seeds of the mutant line were
sterilized for 10 min in a solution of 50% Chlorox with
0.1% SDS, rinsed 3 to 5 times with sterile dH₂O, dried
35 thoroughly on sterile filter paper, and then 2-3 seeds

SECRET 12393 00000000

were sown in liquid B5 medium in 250 mL Belco flasks. The flasks were capped, placed on a rotary shaker at 70-80 rpm, and incubated for 3-4 weeks. Prior to inoculation with Agrobacterium, root tissues were

5 cultured on callus induction medium (MSKig). Roots were harvested by removing the root mass from the Belco flask, placing it in a petri dish, and, using forceps, pulling small bundles of roots from the root mass and placing them on MSKig medium. Petri dishes were sealed

10 with filter tape and incubated for four days.

Agrobacterium strain LBA4404 carrying the plasmids pAL4404 and pAW50 were grown in 5 mL of YEB broth containing 25 mg/L kanamycin and 100 mg/L rifampicin. The culture was grown for approximately 17-20 h in glass

15 culture tubes in a New Brunswick platform shaker (225 rpm) maintained at 28°C. Pre-cultured roots were cut into 0.5 cm segments and placed in a 100 µm filter, made from a Tri-Pour beaker (VWR Scientific, San Francisco, CA USA) and wire mesh, which is set in a petri dish.

20 Root segments were inoculated for several min in 30-50 mL of a 1:20 dilution of the overnight Agrobacterium culture with periodic gentle mixing. Inoculated roots were transferred to sterile filter paper to draw off most of the liquid. Small bundles of roots, consisting

25 of several root segments, were placed on MSKig medium containing 100 µM acetosyringone (3',5'-Dimethoxy-4'-hydroxyaceto-phenone, Aldrich Chemical Co., Milwaukee, WI, USA). Petri plates were sealed with parafilm or filter tape and incubated for 2 to 3 days.

30 After infection, root segments were rinsed and transferred to shoot induction medium with antibiotics. Root bundles were placed in a 100 µm filter unit (described above) and rinsed with 30-50 mL liquid MSKig medium. The filter was vigorously shaken in the

35 solution to help remove the Agrobacterium, transferred

to a clean petri dish, and rinsed again. Roots were blotted on sterile filter paper and bundles of roots were placed on MSg medium containing 500 mg/l vancomycin and either 10 or 20 ppb chlorsulfuron. Plates were
 5 sealed with filter tape and incubated for 12 to 14 days.

Green nodules and small shoot primordia were visible at about 2-3 weeks. The explants were either left intact or were broken into numerous pieces and placed on GM medium containing 200-300 mg/L vancomycin
 10 and either 10 or 20 ppb chlorsulfuron for further shoot development. Plates were either sealed with two pieces of tape or with filter tape. As they developed, individual shoots were isolated from the callus and were placed on MSRg medium containing 100 mg/L vancomycin and
 15 either 10 or 20 ppb chlorsulfuron. Dishes were sealed as described above and incubated for seven to 10 days. Shoots were then transferred to GM medium containing 100-200 mg/L vancomycin in 25x100 petri dishes or Magenta G7 vessels. Many primary transformants (T1)
 20 which were transferred to individual containers set seed (T2).

T2 seed was harvested from selected putative transformants and sown on GM medium containing 10ppb chlorsulfuron. Plates were sealed with filter tape,
 25 cold treated for 2 or more days at 4°C, and then incubated for 10 to 20 days at 23°C under constant illumination as described above. Seedlings were scored as resistant (green, true leaves develop) and sensitive (no true leaves develop).

30 Selected chlorsulfuron resistant T2 seedlings were transplanted to soil and were grown to maturity at 23°C daytime (16 h) 18°C nighttime (8 h) at 65-80% relative humidity.

35 T2 seeds from two plants were harvested at maturity and analysed individually for fatty acid composition by

2025 RELEASE UNDER E.O. 14176

gas chromatography of the fatty acyl methyl esters essentially as described by Browse et al., (Anal. Biochem. (1986) 152:141-145) except that 2.5% H₂SO₄ in methanol was used as the methylation reagent and samples were heated for 1.5 h at 80°C to effect the methanolysis of the seed triglycerides. The results are shown in Table 7.

TABLE 7
Percent Fatty Acid in Seeds of
Transgenic Mutant 3707

Seed Sample	16:0	18:0	18:1	18:2	18:3
wildtype (6)	6	4	14	30	19
mutant 3707 (6)	6	4	14	44	3
1-1	10	4	22	9	55
1-2	11	6	22	14	48
1-3	12	7	16	6	57
1-4	10	4	30	52	4
1-5	10	4	18	17	48
1-6	10	5	15	15	53
2-1	11	5	19	60	4
2-2	10	5	19	9	56
2-3	9	4	27	8	52
2-4	10	5	17	10	56
2-5	10	5	19	9	56
2-6	10	5	17	17	48

The fatty acid composition of the wild-type and mutant line 3707 represents the average of 6 single seeds each. Seeds from plant 1 are designated 1-1 to 1-6 and those from plant 2 are designated 2-1 to 2-6. The 20:1 and 20:2 amounts are not shown. The data shows that the one out of six seeds in each plant show the mutant fatty acid phenotype, while the remaining seeds show more than 10-fold increase in 18:3 to ca.55%. While most of the increase occurs at the expense of 18:2, some of it also occurs at the expense of 18:1.

- Such high levels are of linolenic acid in vegetable oils are observed in specialty oil crops, such as linseed. Thus, overexpression of this gene in other oilscrops, especially canola, which is a close relative of
- 5 Arabidopsis, is also expected to result in such high levels of 18:3.

TABLE 8

Medium Composition

YEP MEDIUM

Bacto Beef Extract	5.0 g
Bacto Yeast Extract	1.0 g
Peptone	5.0 g
Sucrose	5.0 g
MgSO ₄ ·7H ₂ O	0.5 g
Agar (optional)	15.0 g
pH	

VITAMIN SUPPLEMENT

10 mg/L thiamine
50 mg/L pyridoxine
50 mg/L nicotinic acid

MSKIg = Callus Induction Medium

Basic Medium

2% glucose	20 g/L
0.5 mg/L 2,4-D	2.3 µL
0.3 mg/L Kinetin	1.4 µM
5 mg/L IAA	28.5 µM

MSRg = Shoot Induction Medium

Basic Medium

2% glucose	20 g/L
12 mg/L IBA	58.8 µM
0.1 mg/L Kinetin	0.46 µM

BASIC MEDIUM

1 Pkg. Murashige and Skoog Minimal Organics Medium without Sucrose (Gibco #510-3118 or Sigma #M6899)	
10 mL Vitamin Supplement	
0.05% MES	0.5 g/L
0.8% agar	8 g/L
pH	

GM = Germination Medium

Basic Medium

1% sucrose	10 g/L
------------	--------

MSg = Shoot Induction Medium

Basic Medium

2% glucose	20 g/L
0.15 mg/L IAA	0.86 µM
5.0 mg/L 2iP	24.6 µM

EXAMPLE 10

Construction of Vectors for Transformation
of Brassica napus for Reduced Expression
of Delta-15 Desaturases in Developing Seeds

5 Detailed procedures for manipulation of DNA
fragments by restriction endonuclease digestion, size
separation by agarose gel electrophoresis, isolation of
DNA fragments from agarose gels, ligation of DNA
fragments, modification of cut ends of DNA fragments and
10 transformation of E. coli cells with circular DNA
plasmids are all described in Sambrook et al.,
(Molecular Cloning, A Laboratory Manual, 2nd ed (1989)
Cold Spring Harbor Laboratory Press) and Ausubel et al.,
Current Protocols in Molecular Biology (1989) John Wiley
15 & Sons).

Sequences of the cDNA's encoding the B. napus
cytoplasmic delta-15 desaturase and the Brassica napus
plastid delta-15 desaturase were placed in the antisense
orientation behind the promoter region from the a'
20 subunit of the soybean storage protein β -conglycinin to
provide embryo specific expression and high expression
levels.

An embryo-specific expression cassette was
constructed to serve as the basis for chimeric gene
25 constructs for anti-sense expression of the nucleotide
sequences of delta-15 desaturase cDNAs. The vector
pCW109 was produced by the insertion of 555 base pairs
of the β -conglycinin (a' subunit of the 7s seed storage
protein) promoter from soybean (Glycine max), the
30 β -conglycinin 5' untranslated region followed by a
multiple cloning sequence containing the restriction
endonuclease sites for Nco I, Sma I, Kpn I and Xba I,
then 1174 base pairs of the common bean phaseolin 3'
untranslated region into the Hind III site in the
35 cloning vector pUC18 (BRL). The β -conglycinin promoter

5

10

15

20

25

30

The transcription unit [β -conglycinin

An antisense construction based on the plastid delta-15 desaturase was made with the 425 most 3' bases of SEQ ID NO:8 which is contained in the plasmid pBNSFD-8. pBNSFD-8 represents a cDNA of the plastid delta-15 desaturase in pBluescript. The cDNA insert was removed from pBNSFD-8 by digestion with Xho I and Sma I, the fragments were blunted, and the 425 base insert isolated by gel purification. The isolated fragment was cloned into the Sma I site of pCW109A and the antisense orientation of the chosen clone confirmed by digestion of the plasmid with Pst I. Pst I cuts in the plastid delta-15 sequence and in the pCW109A vector 5' to the β -conglycinin promoter to release a 1.2 kB fragment indicative of the antisense orientation. The plasmid containing this construction was called pCCdFdR1.

Digestion of pCCdFdR1 with Hind III removes a 2.3
kB fragment containing the transcriptional unit
[β -conglycinin promoter:plastid delta-15 antisense:3'-
phaseolin sequence]. The fragment was gel isolated and
35 cloned into Hind III digested pBluescript. The

orientation of the fragment was relative to the Bam HI site in the cloning region of pBluescript was determined by digestion with Pst. I as described above. A clone oriented with the promoter toward the Sal I containing end was chosen and given the name pCCdFdR2.

pCCdFdR2 was digested with Bam HI and Sal I, the released fragment was gel isolated and ligated into pZ199 which had been digested with Bam HI and Sal I to give the binary vector pZCCdFdR.

Vectors for transformation of the antisense delta-15 desaturase constructions under control of the β -conglycinin promoter into plants using Agrobacterium tumefaciens were produced by constructing a binary Ti plasmid vector system (Bevan, (1984) Nucl. Acids Res. 12:8711-8720). The starting vector used for these systems (pZS199) is based on a vector which contains: (1) the chimeric gene nopaline synthase/neomycin phosphotransferase as a selectable marker for transformed plant cells (Bevan et al., (1984) Nature 304:184-186), (2) the left and right borders of the T-DNA of the Ti plasmid (Bevan et al., (1984) Nucl. Acids Res. 12:8711-8720), (3) the E. coli lacZ α -complementing segment (Vieria and Messing (1982) Gene 19:259-267) with unique restriction endonuclease sites for Eco RI, Kpn I, Bam HI, Hin DIII, and Sal I, (4) the bacterial replication origin from the Pseudomonas plasmid pVS1 (Itoh et al., (1984) Plasmid 11:206-220), and (5) the bacterial neomycin phosphotransferase gene from Tn5 (Berg et al., (1975) Proc. Natnl. Acad. Sci. U.S.A. 72:3628-3632) as a selectable marker for transformed A. tumefaciens. The nopaline synthase promoter in the plant selectable marker was replaced by the 35S promoter (Odell et al. (1985) Nature, 313:810-813) by a standard restriction endonuclease digestion and ligation strategy. The 35S promoter is

required for efficient Brassica napus transformation as described below.

EXAMPLE 11

AGROBACTERIUM MEDIATED TRANSFORMATION

5 OF BRASSICA NAPUS

The binary vectors pZCC3FdR abd pZCCdFdR were transferred by a freeze/thaw method (Holsters et al., (1978) Mol Gen Genet 163:181-187) to the Agrobacterium strain LBA4404/pAL4404 (Hoekema et al., (1983), Nature
10 303:179-180).

Brassica napus cultivar "Westar" was transformed by co-cultivation of seedling pieces with disarmed Agrobacterium tumefaciens strain LBA4404 carrying the the appropriate binary vector.

15 B. napus seeds were sterilized by stirring in 10% Chlorox, 0.1% SDS for thirty min, and then rinsed thoroughly with sterile distilled water. The seeds were germinated on sterile medium containing 30 mM CaCl₂ and 1.5% agar, and grown for six days in the dark at 24°C.

20 Liquid cultures of Agrobacterium for plant transformation were grown overnight at 28°C in Minimal A medium containing 100 mg/L kanamycin. The bacterial cells are pelleted by centrifugation and resuspended at a concentration of 10⁸ cells/mL in liquid Murashige and
25 Skoog Minimal Organic medium containing 100 µM acetosyringone.

B. napus seedling hypocotyls were cut into 5 mm segments which were immediately placed into the bacterial suspension. After 30 min, the hypocotyl
30 pieces were removed from the bacterial suspension and placed onto BC-12 callus medium containing 100 µM acetosyringone. The plant tissue and Agrobacteria were co-cultivated for three days at 24°C in dim light.

The co-cultivation was terminated by transferring
35 the hypocotyl pieces to BC-12 callus medium containing

862227 52557260

200 mg/L carbenicillin to kill the Agrobacteria, and 25 mg/L kanamycin to select for transformed plant cell growth. The seedling pieces were incubated on this medium for three weeks at 24°C under continuous light.

5 After three weeks, the segments were transferred to BS-48 regeneration medium containing 200 mg/L carbenicillin and 25 mg/L kanamycin. Plant tissue was subcultured every two weeks onto fresh selective regeneration medium, under the same culture conditions
10 described for the callus medium. Putatively transformed calli grow rapidly on regeneration medium; as calli reached a diameter of about 2 mm, they were removed from the hypocotyl pieces and placed on the same medium lacking kanamycin.

15 Shoots began to appear within several weeks after transfer to BS-48 regeneration medium. As soon as the shoots formed discernable stems, they were excised from the calli, transferred to MSV-1A elongation medium, and moved to a 16:8 h day/night photoperiod at 24°C.

20 Once shoots had elongated several internodes, they were cut above the agar surface and the cut ends were dipped in Rootone. Treated shoots were planted directly into wet Metro-Mix 350 soilless potting medium. The pots were covered with plastic bags which were removed when
25 the plants were clearly growing -- after about 10 days.

Plants were grown under a 16:8 h day/night photoperiod, with a daytime temperature of 23°C and a nighttime temperature of 17°C. When the primary flowering stem began to elongate, it was covered with a
30 mesh pollen-containment bag to prevent outcrossing. Self-pollination was facilitated by shaking the plants several times each day. Seeds derived from self-pollinations were harvested about three months after planting.

BBE22T SE6T560

TABLE 9

Minimal A Bacterial Growth Medium	Brassica Callus Medium BC-12
Dissolve in distilled water:	Per liter:
10.5 g potassium phosphate, dibasic	Murashige and Skoog Minimal
4.5 g potassium phosphate, monobasic	Organic Medium (MS salts, 100 mg/L i-inositol, 0.4 mg/L thiamine; GIBCO #510-3118)
1.0 g ammonium sulfate	30 sucrose
0.5 g sodium citrate, dihydrate	18 g mannitol
Make up to 979 mLs with distilled water	1.0 mg/L 2,4-D
Autoclave	3.0 mg/L kinetin
Add 20 mLs filter-sterilized 10% sucrose	0.6% agarose
Add 1 mL filter-sterilized 1 M MgSO ₄	pH 5.8
Brassica Regeneration Medium BS-48	Brassica Shoot Elongation
Murashige and Skoog Minimal	Medium MSV-1A
Organic Medium Gamborg B5 Vitamins (SIGMA #1019)	Murashige and Skoog Minimal
10 g glucose	Organic Medium Gamborg B5
250 mg xylose	Vitamins
600 mg MES	10 g sucrose
0.4% agarose	0.6% agarose
pH 5.7	pH 5.8
Filter-sterilize and add after autoclaving:	
2.0 mg/L zeatin	
0.1 mg/L IAA	

EXAMPLE 12ANALYSIS OF TRANSGENIC BRASSICA NAPUS PLANTS

5 Insertion of the intact antisense transcriptional unit was verified by Southern analysis using transgenic plant leaf tissue as the source of DNA as described in Example 5. Ten micrograms of leaf DNA was digested to completion with a mixture of Bam HI and Sal I

restriction endonucleases and then separated by agarose gel electrophoresis. The separated DNA was transferred to Hybond H⁺ membrane and hybridized with radiolabeled insert from pBNSF3-2. An estimate of the number of

5 copies of the inserted transgene was made by calibrating each Southern blot with standard amounts of pBNSF3-2 corresponding to 1 and 5 copies per genome and comparing intensities of the autoradiographic signal from the

10 standards, the endogenous delta-15 desaturase signals and the inserted gene signal. To date, 38 independent transformants have been analyzed for presence of the gene and 36 were found to be positive.

The relative content of the 5 most abundant fatty acids in canola seeds was determined either by direct

15 trans-esterification of individual seeds in 0.5 mL of methanolic H₂SO₄ (2.5%) or by hexane extraction of bulk seed samples followed by trans-esterification of an aliquot in 0.8 mL of 1% sodium methoxide in methanol. Fatty acid methyl esters were extracted from the

20 methanolic solutions into hexane after the addition of an equal volume of water.

The relative content of 18:3 fatty acid varies significantly during seed development. To a lesser extent, the ratio of 18:3 to 18:2 varies also. Thus

25 meaningful data can be obtained only from seeds after maturation and drydown. Additionally, the ratio of 18:3 to total fatty acid content and to 18:0 varies significantly due to environmental factors, primarily temperature. In this circumstance, the most appropriate

30 controls are the transformed plants which by Southern analysis do not contain the antisense delta-15 transgene. Analysis from the first 5 transformants to reach dry seed are given in Table 10 below. Seeds were harvested using a hand thresher, bulked and a 1.5 g

35 (about 300 seeds) sample was taken. Seed from each

transformant was crushed with a mortar and pestel, extracted 4 times with 8 mL hexane at about 50°C. The combined extracts were reduced in volume to 5 mL and two 50 microliter aliquots were taken for esterification as described above. Separation of the fatty acid methyl esters was done by gas-liquid chromatography using an Omegawax 320 column (Supelco Inc., 0.32 mm ID X 30M) run isothermally at 220° and cycled to 260° between each injection.

TABLE 10

<u>Transformant</u> <u>No.</u>	<u>% 18:3</u>	<u>%18:3/18:2</u>	<u>Antisense</u> <u>delta-15</u> <u>Copy No.</u>
pZCC3FdR-91	6.2	0.39	0
pZCC3FdR-81	5.9	0.33	1
pZCC3FdR-15	6.0	0.38	2
pZCC3FdR-11	5.6	0.34	1
pZCC3FdR-148	8.2	0.40	2

The differences between the 4 transformed lines and line 92 are very small, however to test the significance of the difference in the 18:3/18:2 ratio between line 81 and 91, 25 individual seeds from each line were transesterified and their fatty acid composition determined. The average ratio for line 81 was 0.345 with a coefficient of variation of 11.6% while the average for line 91 was 0.375 with a coefficient of variation of 8.0%. The sample means are significantly different at the 0.01% level using Student's t test.

EXAMPLE 13

CONSTRUCTION OF VECTORS FOR TRANSFORMATION OF
GLYCINE MAX FOR REDUCED EXPRESSION OF DELTA-15
DESATURASES IN DEVELOPING SEEDS

The antisense G. max plastid delta-15 desaturase cDNA sequence under control of the β -conglycinin promoter was constructed using the vector pCW109A described in Example 10 above. For use in the soybean

transformation system described below, the transcriptional unit was placed in a vector along with an appropriate selectable marker expression system. The starting vector was pML45, which consists of the non-tissue specific and constitutive promoter designated 508D and described in Hershey (WO 9011361) driving expression of the neomycin phosphotransferase gene described in (Beck et al. (1982) Gene 19:327-336) followed by the 3' end of the nopaline synthase gene including nucleotides 848 to 1550 described by (Depicker et al. (1982) J. Appl. Genet. 1:561-574). This transcriptional unit was inserted into the commercial cloning vector pGEM9Z (BRL) and is flanked at the 5' end of the 508D promoter by the restriction sites Sal I, Xba I, Bam HI and Sma I in that order. An additional Sal I site is present at the 3' end of the NOS 3' sequence and the Xba I, Bam HI and Sal I sites are unique.

Removal of the unit[β -conglycinin promoter:cloning region:phaseolin 3' end] from pCW109A by digestion with Hind III, blunting the ends and isolating the 1.8 kB fragment afforded the expression cassette pCST by ligating the above isolated fragment into the Sma I site of pML45. A clone with the β -conglycinin promoter in the same orientation as the 508D promoter were chosen by digestion with Xba I. The correct orientation releases a 700 bp fragment. This vector cassette was called pCST.

The 2.2 kB insert encoding the soybean, plastid delta-15 desaturase was subcloned from the plasmid pXF1 by digestion with HinP I to remove about 1 kB of unrelated cDNA. HinP I cuts within the cDNA insert very near the 5' end of the cDNA for the delta-15 desaturase and about 300 bp from the 3' end of that cDNA. The Cla I compatible ends were cloned into Cla I digested pBluescript and a clone with the 5' end of the cDNA

toward the Eco RV site in the pBluescript cloning region was selected based on the release of a 900 bp fragment by digestion with Pst I. The subcloned plasmid was called pS3Fd1.

- 5 The delta-15 encoding sequence was removed from pS3Fd1 by digestion with Hinc II and Eco RV, the 2.2 kB fragment was gel isolated and cloned into the opened Sma I site in pCST1. A clone with the delta-15 sequence in the antisense orientation to the β -conglycinin promoter
- 10 was selected by digestion with Xba I. The antisense construct releases a 400 bp piece and that clone was designated pCS3FdST1R.

EXAMPLE 14

TRANSFORMATION OF SOMATIC SOYBEAN EMBRYO CULTURES

- 15 Soybean embryogenic suspension cultures are maintained in 35 mL liquid media (SB55 or SBP6) on a rotary shaker, 150 rpm, at 28°C with mixed florescent and incandescent lights on a 16:8 h day/night schedule. Cultures were subcultured every four weeks by
- 20 inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

- Soybean embryogenic suspension cultures were transformed with pCS3FdST1R by the method of particle gun bombardment (see Kline et al. (1987) Nature (London)
- 25 327:70). A Du Pont Biolistic PDS1000/HE instrument (helium retrofit) was used for these transformations.

- To 50 mL of a 60 mg/mL 1 mm gold particle suspension was added (in order); 5 μ L DNA (1 μ g/ μ L), 20 μ L spermidine (0.1M), and 50 μ L CaCl₂ (2.5 M). The
- 30 particle preparation was agitated for 3 min, spun in a microfuge for 10 sec and the supernatant removed. The DNA-coated particles were then washed once in 400 μ L 70% ethanol and resuspended in 40 μ L of anhydrous ethanol. The DNA/particle suspension was sonicated three times

for 1 sec each. Five μ L of the DNA-coated gold particles were then loaded on each macro carrier disk.

Approximately 300-400 mg of a four week old suspension culture was placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue were normally bombarded. Membrane rupture pressure was set at 1000 psi and the chamber was evacuated to a vacuum of 28 inches of mercury. The tissue was placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue was placed back into liquid and cultured as described above.

Eleven days post bombardment, the liquid media was exchanged with fresh SB55 containing 50 mg/mL hygromycin. The selective media was refreshed weekly. Seven weeks post bombardment, green, transformed tissue was observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue was removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Thus each new line was treated as independent transformation event. These suspensions can then be maintained as suspensions of embryos clustered in an immature developmental stage through subculture or regenerated into whole plants by maturation and germination of individual somatic embryos.

Transformed embryogenic clusters were removed from liquid culture and placed on a solid agar media (SB103) containing no hormones or antibiotics. Embryos were cultured for eight weeks at 26°C with mixed florescent and incandescent lights on a 16:8 h day/night schedule. During this period, individual embryos were removed from the clusters and analyzed at various stages of embryo

35221 35221

development After eight weeks the embryos become suitable for germination.

TABLE 11

Media:		B5 Vitimin Stock
SB55 and SBP6 Stock Solutions (g/L):		10 g m-inositol
MS Sulfate 100X Stock		100 mg nicotinic acid
MgSO ₄ 7H ₂ O	37.0	100 mg pyridoxine HCl
MnSO ₄ H ₂ O	1.69	1 g thiamine
ZnSO ₄ 7H ₂ O	0.86	SB55 (per Liter)
CuSO ₄ 5H ₂ O	0.0025	10 mL each MS stocks
MS Halides 100X Stock		1 mL B5 Vitaimin stock
CaCl ₂ 2H ₂ O	44.0	0.8 g NH ₄ NO ₃
KI	0.083	3.033 g KNO ₃
CoCl ₂ 6H ₂ O	0.00125	1 mL 2,4-D (10mg/mL stock)
KH ₂ PO ₄	17.0	60 g sucrose
H ₃ BO ₃	0.62	0.667 g asparagine
Na ₂ MoO ₄ 2H ₂ O	0.025	pH 5.7
MS FeEDTA 100X Stock		For SBP6- substitute 0.5 mL 2,4-D
Na ₂ EDTA	3.724	SB103 (per Liter)
FeSO ₄ 7H ₂ O	2.784	MS Salts
		6% maltose
		750 mg MgCl ₂
		0.2% Gelrite
		pH 5.7

EXAMPLE 15

ANALYSIS OF TRANSGENIC GLYCINE MAX PLANTS

- 5 While in the globular embryo state in liquid culture as described in Example 14, somatic soybean embryos contain very low amounts of triacylglycerol or storage proteins typical of maturing, zygotic soybean embryos. At this developmental stage, the ratio of
- 10 total triacylglyceride to total polar lipid (phospholipids and glycolipid) is about 1:4, as is typical of zygotic soybean embryos at the developmental

stage from which the somatic embryo culture was initiated. At the globular stage as well, the mRNAs for the prominent seed proteins (α' subunit of β -conglycinin, Kunitz Trypsin Inhibitor III and Soybean Seed Lectin) are essentially absent. Upon transfer to hormone free media to allow differentiation to the maturing somatic embryo state as described in Example 14, triacylglycerol becomes the most abundant lipid class. As well, mRNAs for α' -subunit of β -conglycinin, Kunitz Trypsin Inhibitor III and Soybean Seed Lectin become very abundant messages in the total mRNA population. In these respects the somatic soybean embryo system behaves very similarly to maturing zygotic soybean embryos *in vivo*, and is therefore a good and rapid model system for analyzing the phenotypic effects of modifying the expression of genes in the fatty acid biosynthesis pathway. Similar somatic embryo culture systems have been documented and used in another oilseed crop, rapeseed (Taylor et al. (1990) *Planta* 181:18-26). Fatty acid analysis was performed as described in Example 12 using single embryos as the tissue source. A number of embryos from line 2872 (control tissue transformed with pCST) and lines 299,303,306 and 307 (line 2872 transformed with plasmid pCS3FdST1R) were analyzed for fatty acid content. The relative fatty-acid composition of embryos taken from tissue transformed with pCS3FdST1R was compared with control tissue, transformed with pCST. The results of this analysis are shown in Table 12.

TABLE 12

<u>Line</u>	<u>Embryo</u>	<u>16:0</u>	<u>18:0</u>	<u>18:1</u>	<u>18:2</u>	<u>18:3</u>
2872	1	17.7	4.1	11.3	52.8	14.1
	2	17.3	4.3	10.9	49.5	18.0
	3	16.1	4.1	13.8	48.2	17.3
	4	17.5	3.6	11.7	52.0	14.1

<u>Line</u>	<u>Embryo</u>	<u>16:0</u>	<u>18:0</u>	<u>18:1</u>	<u>18:2</u>	<u>18:3</u>
	5	16.6	3.9	12.7	53.7	12.6
	6	14.8	3.0	14.7	55.3	11.1
	av	16.7	3.8	12.5	51.9	14.5
299-1-3	1	16.5	4.1	9.7	61.4	6.3
299-15-1	1	14.7	3.6	11.9	61.3	8.4
	2	16.6	3.7	12.1	58.6	8.6
	3	16.7	4.1	14.9	53.2	11.1
	4	15.2	4.0	9.1	60.2	11.5
	5	16.0	4.2	13.9	55.2	10.7
	6	15.2	3.5	9.9	63.4	8.1
303-7-1	1	14.1	2.2	10.6	59.4	13.7
	2	14.0	2.8	12.5	59.3	11.4
306-4-5	1	17.5	4.2	8.1	62.7	7.4
	215.7	3.3	9.0	60.5	11.5	
	3	17.1	3.4	9.3	60.7	9.5
	4	15.7	3.8	9.2	61.2	9.7
	5	17.7	3.9	6.5	58.3	13.6
	6	16.6	3.4	10.2	59.2	10.6
306-4-8	1	16.6	3.9	15.3	50.7	11.8
	2	17.8	3.6	15.7	50.0	10.8
	3	16.7	3.3	11.1	52.0	14.6
	4	19.0	4.0	10.3	53.1	12.3
	5	19.7	3.5	9.0	53.6	13.0
	6	18.0	2.9	13.1	52.8	10.9
307-1-1	1	14.4	3.7	11.2	64.4	6.3
	2	15.4	3.4	7.8	61.0	11.3
	3	17.2	2.5	12.0	57.2	11.1
307-1-2	1	13.4	3.0	8.4	55.4	19.9
	2	16.3	3.1	6.4	55.7	18.7
	3	14.0	3.3	8.8	58.7	15.2
	4	15.8	2.5	9.8	59.7	12.2
	5	14.6	3.7	14.9	51.1	15.7
	6	14.3	3.9	11.4	55.5	14.1

862227 526746

<u>Line</u>	<u>Embryo</u>	<u>16:0</u>	<u>18:0</u>	<u>18:1</u>	<u>18:2</u>	<u>18:3</u>
307-1-3	1	14.8	3.1	9.4	60.5	12.2
	2	18.0	3.0	5.3	56.2	15.2
	3	18.0	3.4	2.5	58.6	15.4
307-1-4	1	15.0	2.7	13.8	61.7	6.9
	2	15.9	2.7	9.8	62.0	9.6
	3	14.6	3.2	13.4	61.4	6.7
307-1-5	1	15.9	3.5	7.6	61.7	11.2
	2	14.6	3.5	10.0	61.3	10.6
	3	18.7	2.6	6.8	53.0	19.0
307-1-7	1	15.3	3.5	12.5	60.3	8.5
	2	16.2	2.2	13.9	57.1	10.6
	3	14.9	3.1	12.2	58.0	11.8
307-1-9	1	16.4	2.9	23.2	47.9	9.6
	2	19.6	0.0	20.4	51.3	8.8
	3	16.8	3.3	24.6	49.6	5.7
307-1-11	1	18.1	3.6	5.7	52.9	19.7
	2	14.7	3.7	9.9	58.7	13.0
	3	15.1	3.7	11.3	55.8	14.1

The average 18:3 content of control embryos was 14.5% with a range from 11.1% to 18.0%. The average 18:3 content of transformed embryos was 11.5% with a range of 6.3% to 19.9%. Almost 80% of the transformed embryos (38/48) had an 18:3 content below that of the control mean. About 44% had an 18:3 content less than the lowest observed control value and 12.5% had an 18:3 content less than half of the control mean value (i.e., less than 7.5%). The lowest 18:3 content observed in transformed tissue was 6.3% (299-1-3, 307-1-2 #1) compared with the control low of 11.1%. In all cases in transformed tissue, a decrease in 18:3 content was reflected by an equivalent increase in 18:2 content indicating that the desaturation of 18:2 to 18:3 had

been reduced. The relative content of the the other fatty acids remained unchanged.

Southern analysis for the presence of the intact, introduced antisense construction was performed, as described in Example 12 using Bam HI cut gDNA, on a number of the transformed lines listed below using groups of embryos from a single transformation event. The approximate intact antisense copy number was estimated from the number and intensity of hybridizing bands on the autoradiograms and is shown in Table 13.

TABLE 13

<u>Line No.</u>	<u>Antisense copy No.</u>	<u>18:3 (low)</u>	<u>18:3 (average)</u>	<u>18:2/18:3 ratio</u>
2872	0	11.1	14.5	3.6
303-7/1	1	11.4	12.6	4.7
307-1/2	3	12.2	16.0	3.5
306-4/8	3	10.8	12.2	4.3
307-1/7	4	8.5	10.3	5.7
306-4/5	6	7.4	10.4	5.8
307-1/1	6	6.3	9.6	6.3
299-15/1	7	8.1	9.7	6.1
307-1/4	8	6.7	7.7	8.0

There was a reasonable correlation between intact antisense copy number and 18:3 content, an increase in copy number correlating with a decreased 18:3 content and a consequent increase in the 18:2/18:3 ratio. The average 18:2/18:3 ratio of line 307-1/4, which had at least 8 copies of the antisense cDNA, was more than twice that of the control.